

**C1orf96 antibody - middle region**  
**Rabbit Polyclonal Antibody**  
**Catalog # AI13465****Specification**

---

**C1orf96 antibody - middle region - Product Information**

|                   |                                                                |
|-------------------|----------------------------------------------------------------|
| Application       | WB                                                             |
| Primary Accession | <a href="#">Q6IQ19</a>                                         |
| Other Accession   | <a href="#">NM_145257</a> , <a href="#">NP_660300</a>          |
| Reactivity        | Human, Mouse, Rat, Rabbit, Pig, Horse, Bovine, Guinea Pig, Dog |
| Predicted         | Rabbit, Chicken, Horse, Bovine, Guinea Pig, Dog                |
| Host              | Rabbit                                                         |
| Clonality         | Polyclonal                                                     |
| Calculated MW     | 30 KDa                                                         |

**C1orf96 antibody - middle region - Additional Information****Gene ID** 126731**Alias Symbol** **FLJ37296, FLJ41471, RP4-803J11.3, C1orf96**  
**Other Names**  
Centriole, cilia and spindle-associated protein, CCSAP, C1orf96, CSAP**Format**

Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.

**Reconstitution & Storage**

Add 50 ul of distilled water. Final anti-C1orf96 antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.

**Precautions**

C1orf96 antibody - middle region is for research use only and not for use in diagnostic or therapeutic procedures.

**C1orf96 antibody - middle region - Protein Information****Name** CCSAP**Synonyms** C1orf96, CSAP**Function**Plays a role in microtubule (MT) stabilization and this stabilization involves the maintenance of NUMA1 at the spindle poles. Colocalizes with polyglutamylated MTs to promote MT stabilization and regulate bipolar spindle formation in mitosis. Binding of CCSAP to centrosomes and the spindle around centrosomes during mitosis inhibits MT depolymerization, thereby stabilizing the mitotic spindle (PubMed: <http://www.uniprot.org/citations/26562023>)

target="\_blank">26562023</a>). May play a role in embryonic development. May be required for proper cilia beating (By similarity).

#### **Cellular Location**


Cytoplasm, cytoskeleton, microtubule organizing center, centrosome, centriole. Cytoplasm, cytoskeleton, spindle. Cytoplasm, cytoskeleton. Cytoplasm, cytoskeleton, cilium basal body. Cytoplasm, cytoskeleton, cilium axoneme. Cell projection, axon. Cell projection, cilium. Cytoplasm, cytoskeleton, microtubule organizing center, centrosome. Note=Localizes to two to four centrioles throughout the cell cycle. Localizes to mitotic spindle microtubules during prometaphase and throughout the remainder of mitosis. Localizes to cytoskeleton on interphase. Localizes at the ciliary transition zone which connects the basal bodies to ciliary microtubules. Colocalizes with polyglutamylated tubulin.

#### **C1orf96 antibody - middle region - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

#### **C1orf96 antibody - middle region - Images**



WB Suggested Anti-C1orf96 Antibody Titration: 0.2-1 µg/ml

ELISA Titer: 1:312500

Positive Control: Human heart

#### **C1orf96 antibody - middle region - References**

Ota T., et al. Nat. Genet. 36:40-45(2004).  
Gregory S.G., et al. Nature 441:315-321(2006).  
Mural R.J., et al. Submitted (JUL-2005) to the EMBL/GenBank/DDBJ databases.  
Backer C.B., et al. Mol. Biol. Cell 23:2122-2130(2012).  
Van Damme P., et al. Proc. Natl. Acad. Sci. U.S.A. 109:12449-12454(2012).