

C1orf93 antibody - middle region
Rabbit Polyclonal Antibody
Catalog # AI13482**Specification**

C1orf93 antibody - middle region - Product Information

Application	WB
Primary Accession	Q5R7S9
Other Accession	NM_152371 , NP_689584
Reactivity	Human, Mouse, Rat, Pig, Horse, Bovine, Guinea Pig, Dog
Predicted Host	Horse
Clonality	Rabbit
Calculated MW	Polyclonal 21kDa KDa

C1orf93 antibody - middle region - Additional Information**Gene ID** 100190803**Alias Symbol** DKFZp547M123, MGC26818, RP3-395M20.10, C1orf93**Other Names**

Prostamide/prostaglandin F synthase, Prostamide/PG F synthase, Prostamide/PGF synthase, 1.11.1.20, FAM213B

Format

Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.

Reconstitution & Storage

Add 50 ul of distilled water. Final anti-C1orf93 antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.

Precautions

C1orf93 antibody - middle region is for research use only and not for use in diagnostic or therapeutic procedures.

C1orf93 antibody - middle region - Protein Information**Name** PRXL2B**Synonyms** FAM213B**Function**

Catalyzes the reduction of prostaglandin-ethanolamide H(2) (prostamide H(2)) to prostamide F(2alpha) with NADPH as proton donor. Also able to reduce prostaglandin H(2) to prostaglandin F(2alpha) (By similarity).


Cellular Location

Cytoplasm, cytosol {ECO:0000250|UniProtKB:Q9DB60}

C1orf93 antibody - middle region - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

C1orf93 antibody - middle region - Images

WB Suggested Anti-C1orf93 Antibody Titration: 0.2-1 μ g/ml
Positive Control: PANC1 cell lysate