

C12orf50 antibody - N-terminal region
Rabbit Polyclonal Antibody
Catalog # AI13499**Specification**

C12orf50 antibody - N-terminal region - Product Information

Application	WB
Primary Accession	Q8NA57
Other Accession	NM_152589 , NP_689802
Reactivity	Human, Rat, Rabbit, Pig, Bovine, Guinea Pig, Dog
Predicted Host	Human, Rabbit, Pig, Guinea Pig, Dog
Clonality	Rabbit
Calculated MW	Polyclonal 47kDa KDa

C12orf50 antibody - N-terminal region - Additional Information**Gene ID** 160419

Alias Symbol	FLJ35821
Other Names	
Uncharacterized protein C12orf50, C12orf50	

Format

Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.

Reconstitution & Storage

Add 50 ul of distilled water. Final anti-C12orf50 antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.

Precautions

C12orf50 antibody - N-terminal region is for research use only and not for use in diagnostic or therapeutic procedures.


C12orf50 antibody - N-terminal region - Protein Information**Name** C12orf50**C12orf50 antibody - N-terminal region - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)

- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

C12orf50 antibody - N-terminal region - Images



WB Suggested Anti-C12orf50 Antibody Titration: 0.2-1 μ g/ml

ELISA Titer: 1:62500

Positive Control: HepG2 cell lysate

C12orf50 antibody - N-terminal region - References

Ota T., et al. Nat. Genet. 36:40-45(2004).