

Coil Antibody - C-terminal region
Rabbit Polyclonal Antibody
Catalog # AI13587**Specification**

Coil Antibody - C-terminal region - Product Information

Application	WB
Primary Accession	O923T0
Other Accession	NM_017360 , NP_059056
Reactivity	Human, Mouse, Rat, Rabbit, Pig, Horse, Bovine, Guinea Pig, Dog
Predicted	Human, Mouse, Rat, Rabbit, Pig, Horse, Bovine, Guinea Pig, Dog
Host	Rabbit
Clonality	Polyclonal
Calculated MW	62kDa KDa

Coil Antibody - C-terminal region - Additional Information**Format**

Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.

Reconstitution & Storage

Add 50 ul of distilled water. Final anti-Coil antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.

Precautions


Coil Antibody - C-terminal region is for research use only and not for use in diagnostic or therapeutic procedures.

Coil Antibody - C-terminal region - Protein Information**Coil Antibody - C-terminal region - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Coil Antibody - C-terminal region - Images



Host: Rabbit
Target Name: Coil
Sample Tissue: Rat Testis lysates
Antibody Dilution: 1.0 μ g/ml