

**CCT6B antibody - N-terminal region**  
**Rabbit Polyclonal Antibody**  
**Catalog # AI13596****Specification**

---

**CCT6B antibody - N-terminal region - Product Information**

|                   |                                                           |
|-------------------|-----------------------------------------------------------|
| Application       | WB                                                        |
| Primary Accession | <a href="#">O92526</a>                                    |
| Other Accession   | <a href="#">NM_006584</a> , <a href="#">NP_006575</a>     |
| Reactivity        | Human, Mouse, Rat, Rabbit, Horse, Bovine, Guinea Pig, Dog |
| Predicted         | Human, Mouse, Rat, Rabbit, Bovine, Guinea Pig, Dog        |
| Host              | Rabbit                                                    |
| Clonality         | Polyclonal                                                |
| Calculated MW     | 58kDa kDa                                                 |

**CCT6B antibody - N-terminal region - Additional Information****Gene ID** 10693**Alias Symbol** CCT-zeta-2, CCTZ-2, Cctz2, TCP-1-zeta-2, TSA303**Other Names**

T-complex protein 1 subunit zeta-2, TCP-1-zeta-2, CCT-zeta-2, CCT-zeta-like, TCP-1-zeta-like, Testis-specific Tcp20, Testis-specific protein TSA303, CCT6B

**Format**

Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.

**Reconstitution & Storage**

Add 50 ul of distilled water. Final anti-CCT6B antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.

**Precautions**

CCT6B antibody - N-terminal region is for research use only and not for use in diagnostic or therapeutic procedures.

**CCT6B antibody - N-terminal region - Protein Information****Name** CCT6B**Function**

Component of the chaperonin-containing T-complex (TRiC), a molecular chaperone complex that assists the folding of proteins upon ATP hydrolysis.

**Cellular Location**

Cytoplasm.


**Tissue Location**

Testis-specific..

**CCT6B antibody - N-terminal region - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

**CCT6B antibody - N-terminal region - Images**

WB Suggested Anti-CCT6B Antibody Titration: 0.2-1 µg/ml  
Positive Control: 721\_B cell lysate

**CCT6B antibody - N-terminal region - References**

Ozaki K., et al. Genomics 36:316-319(1996).  
Ota T., et al. Nat. Genet. 36:40-45(2004).  
Zody M.C., et al. Nature 440:1045-1049(2006).  
Burkard T.R., et al. BMC Syst. Biol. 5:17-17(2011).