

CCDC70 antibody - middle region
Rabbit Polyclonal Antibody
Catalog # AI13621**Specification**

CCDC70 antibody - middle region - Product Information

Application	WB
Primary Accession	Q6NSX1
Other Accession	NM_031290 , NP_112580
Reactivity	Human, Mouse, Rat, Pig, Horse, Dog
Predicted	Mouse, Pig, Horse, Dog
Host	Rabbit
Clonality	Polyclonal
Calculated MW	26kDa KDa

CCDC70 antibody - middle region - Additional Information**Gene ID** 83446**Alias Symbol** **DKFZP434K1172, FLJ25853****Other Names**

Coiled-coil domain-containing protein 70, CCDC70

Format

Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.

Reconstitution & Storage

Add 50 ul of distilled water. Final anti-CCDC70 antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.

Precautions

CCDC70 antibody - middle region is for research use only and not for use in diagnostic or therapeutic procedures.


CCDC70 antibody - middle region - Protein Information**Name** CCDC70 ([HGNC:25303](#))**CCDC70 antibody - middle region - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)

- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

CCDC70 antibody - middle region - Images



WB Suggested Anti-CCDC70 Antibody Titration: 0.2-1 µg/ml

ELISA Titer: 1:312500

Positive Control: Human Stomach

CCDC70 antibody - middle region - References

Wiemann S., et al. Genome Res. 11:422-435(2001).

Ota T., et al. Nat. Genet. 36:40-45(2004).

Dunham A., et al. Nature 428:522-528(2004).