

C5orf39 antibody - N-terminal region
Rabbit Polyclonal Antibody
Catalog # AI13647**Specification**

C5orf39 antibody - N-terminal region - Product Information

Application	WB
Primary Accession	O3ZCQ2
Other Accession	NM_001014279 , NP_001014301
Reactivity	Human
Predicted	Human
Host	Rabbit
Clonality	Polyclonal
Calculated MW	22kDa KDa

C5orf39 antibody - N-terminal region - Additional Information**Gene ID** 389289**Alias Symbol** AX2R, AXIIR, C5orf39**Other Names**

Annexin-2 receptor, Annexin II receptor, AXIIR, ANXA2R, AX2R, C5orf39

Format

Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.

Reconstitution & Storage

Add 50 ul of distilled water. Final anti-C5orf39 antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.

Precautions

C5orf39 antibody - N-terminal region is for research use only and not for use in diagnostic or therapeutic procedures.

C5orf39 antibody - N-terminal region - Protein Information**Name** ANXA2R**Synonyms** AX2R, C5orf39**Function**

May act as a receptor for annexin II on marrow stromal cells to induce osteoclast formation.

Tissue Location


Widely expressed. Highly expressed in lymphocytes. Expressed in both resting CD4(+) and CD8(+) T-cells

C5orf39 antibody - N-terminal region - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

C5orf39 antibody - N-terminal region - Images



WB Suggested Anti-C5orf39 Antibody Titration: 0.2-1 μ g/ml
Positive Control: Human Placenta

C5orf39 antibody - N-terminal region - References

Lu G., et al. J. Biol. Chem. 281:30542-30550(2006).
Schmutz J., et al. Nature 431:268-274(2004).
Menea C., et al. J. Clin. Invest. 103:1605-1613(1999).