

WDTC1 antibody - N-terminal region
Rabbit Polyclonal Antibody
Catalog # AI13751**Specification**

WDTC1 antibody - N-terminal region - Product Information

Application	WB
Primary Accession	Q5SSC5
Other Accession	NM_015023 , NP_055838
Reactivity	Human, Mouse, Rat, Rabbit, Pig, Horse, Guinea Pig, Dog
Predicted	Human, Mouse, Rat, Rabbit, Pig, Chicken, Horse, Guinea Pig, Dog
Host	Rabbit
Clonality	Polyclonal
Calculated MW	76kDa KDa

WDTC1 antibody - N-terminal region - Additional Information

Alias Symbol **ADP, KIAA1037, RP11-4K3_A.1, DCAF9**

Format

Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.

Reconstitution & Storage

Add 50 ul of distilled water. Final anti-WDTC1 antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.

Precautions


WDTC1 antibody - N-terminal region is for research use only and not for use in diagnostic or therapeutic procedures.

WDTC1 antibody - N-terminal region - Protein Information**WDTC1 antibody - N-terminal region - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

WDTC1 antibody - N-terminal region - Images



WB Suggested Anti-WDTC1 Antibody Titration: 0.2-1 µg/ml

ELISA Titer: 1:312500

Positive Control: 721_B cell lysate

There is BioGPS gene expression data showing that WDTC1 is expressed in 721_B