

DCBLD1 antibody - N-terminal region
Rabbit Polyclonal Antibody
Catalog # AI13837**Specification**

DCBLD1 antibody - N-terminal region - Product Information

Application	WB
Primary Accession	Q8N8Z6
Other Accession	NM_173674 , NP_775945
Reactivity	Human, Mouse, Rat, Rabbit, Pig, Bovine, Dog, Horse, Guinea Pig
Predicted	Human, Rat, Rabbit, Pig, Chicken, Bovine, Dog, Horse, Guinea Pig
Host	Rabbit
Clonality	Polyclonal
Calculated MW	59kDa kDa

DCBLD1 antibody - N-terminal region - Additional Information**Gene ID** 285761**Alias Symbol** [DKFZp686L21135](#), [MGC46341](#), [dJ94G16.1](#)
Other Names
Discoidin, CUB and LCCL domain-containing protein 1, DCBLD1**Format**

Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.

Reconstitution & Storage

Add 50 ul of distilled water. Final anti-DCBLD1 antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.

Precautions

DCBLD1 antibody - N-terminal region is for research use only and not for use in diagnostic or therapeutic procedures.

DCBLD1 antibody - N-terminal region - Protein Information**Name** DCBLD1**Cellular Location**


Membrane; Single-pass type I membrane protein

DCBLD1 antibody - N-terminal region - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

DCBLD1 antibody - N-terminal region - Images



WB Suggested Anti-DCBLD1 Antibody Titration: 0.2-1 μ g/ml

ELISA Titer: 1:62500

Positive Control: OVCAR-3 cell lysate

DCBLD1 antibody - N-terminal region - References

- Ota T., et al. Nat. Genet. 36:40-45(2004).
Mungall A.J., et al. Nature 425:805-811(2003).
Olsen J.V., et al. Cell 127:635-648(2006).
Olsen J.V., et al. Sci. Signal. 3:RA3-RA3(2010).