

MAPK13 antibody - middle region
Rabbit Polyclonal Antibody
Catalog # AI13942**Specification**

MAPK13 antibody - middle region - Product Information

Application	WB
Primary Accession	O9N272
Other Accession	NM_002754 , NP_002745
Reactivity	Human, Mouse, Rat, Rabbit, Pig, Goat, Sheep, Horse, Yeast, Bovine, Guinea Pig, Dog
Predicted	Human, Mouse, Rat, Rabbit, Pig, Chicken, Goat, Sheep, Bovine, Guinea Pig, Dog
Host	Rabbit
Clonality	Polyclonal
Calculated MW	42kDa KDa

MAPK13 antibody - middle region - Additional Information**Gene ID** 462644**Alias Symbol** MGC99536, PRKM13, SAPK4, p38delta**Other Names**

Mitogen-activated protein kinase 13, MAP kinase 13, MAPK 13, 2.7.11.24, Stress-activated protein kinase 4, MAPK13, SAPK4

Format

Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.

Reconstitution & Storage

Add 50 ul of distilled water. Final anti-MAPK13 antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.

Precautions

MAPK13 antibody - middle region is for research use only and not for use in diagnostic or therapeutic procedures.

MAPK13 antibody - middle region - Protein Information**Name** MAPK13**Synonyms** SAPK4**Function**

Serine/threonine kinase which acts as an essential component of the MAP kinase signal transduction pathway. MAPK13 is one of the four p38 MAPKs which play an important role in the cascades of cellular responses evoked by extracellular stimuli such as pro-inflammatory cytokines


or physical stress leading to direct activation of transcription factors such as ELK1 and ATF2. Accordingly, p38 MAPKs phosphorylate a broad range of proteins and it has been estimated that they may have approximately 200 to 300 substrates each. MAPK13 is one of the less studied p38 MAPK isoforms. Some of the targets are downstream kinases such as MAPKAPK2, which are activated through phosphorylation and further phosphorylate additional targets. Plays a role in the regulation of protein translation by phosphorylating and inactivating EEF2K. Involved in cytoskeletal remodeling through phosphorylation of MAPT and STMN1. Mediates UV irradiation induced up- regulation of the gene expression of CXCL14. Plays an important role in the regulation of epidermal keratinocyte differentiation, apoptosis and skin tumor development. Phosphorylates the transcriptional activator MYB in response to stress which leads to rapid MYB degradation via a proteasome-dependent pathway. MAPK13 also phosphorylates and down- regulates PRKD1 during regulation of insulin secretion in pancreatic beta cells (By similarity).

MAPK13 antibody - middle region - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

MAPK13 antibody - middle region - Images



WB Suggested Anti-MAPK13 Antibody Titration: 0.2-1 µg/ml

ELISA Titer: 1:1562500

Positive Control: Jurkat cell lysate

MAPK13 antibody - middle region - References

Herbison C.E., et al. DNA Seq. 10:229-243(1999).