

ALLC antibody - N-terminal region
Rabbit Polyclonal Antibody
Catalog # AI14081**Specification**

ALLC antibody - N-terminal region - Product Information

Application	WB
Primary Accession	B4DY77
Other Accession	NM_018436 , NP_060906
Reactivity	Human, Rabbit, Horse, Bovine, Dog
Predicted	Human, Horse, Dog
Host	Rabbit
Clonality	Polyclonal
Calculated MW	43kDa KDa

ALLC antibody - N-terminal region - Additional Information

Alias Symbol **ALC**

Format

Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.

Reconstitution & Storage

Add 50 ul of distilled water. Final anti-ALLC antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.

Precautions


ALLC antibody - N-terminal region is for research use only and not for use in diagnostic or therapeutic procedures.

ALLC antibody - N-terminal region - Protein Information**ALLC antibody - N-terminal region - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

ALLC antibody - N-terminal region - Images



WB Suggested Anti-ALLC Antibody Titration: 0.2-1 μ g/ml
Positive Control: Jurkat cell lysate