

Hoxa10 antibody - C-terminal region
Rabbit Polyclonal Antibody
Catalog # AI14256**Specification**

Hoxa10 antibody - C-terminal region - Product Information

Application	WB
Primary Accession	B2RYB5
Other Accession	NM_001129878 , NP_001123350
Reactivity	Human, Mouse, Rat, Rabbit, Pig, Sheep, Horse, Bovine, Guinea Pig, Dog
Predicted	Rat
Host	Rabbit
Clonality	Polyclonal
Calculated MW	34kDa KDa

Hoxa10 antibody - C-terminal region - Additional Information

Alias Symbol **MGC188427, RGD1564605, Hoxa10**

Format

Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.

Reconstitution & Storage

Add 50 ul of distilled water. Final anti-Hoxa10 antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.

Precautions


Hoxa10 antibody - C-terminal region is for research use only and not for use in diagnostic or therapeutic procedures.

Hoxa10 antibody - C-terminal region - Protein Information**Hoxa10 antibody - C-terminal region - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Hoxa10 antibody - C-terminal region - Images



WB Suggested Anti-Hoxa10 Antibody Titration: 1.0 μ g/ml
Positive Control: Rat Muscle