

Ubxn6 antibody - middle region
Rabbit Polyclonal Antibody
Catalog # AI14378**Specification**

Ubxn6 antibody - middle region - Product Information

Application	WB
Primary Accession	Q99PL6
Other Accession	NM_024432 , NP_077752
Reactivity	Human, Mouse, Rat, Pig, Horse, Yeast, Bovine, Guinea Pig, Dog
Predicted	Human, Mouse, Chicken, Horse, Bovine, Guinea Pig, Dog
Host	Rabbit
Clonality	Polyclonal
Calculated MW	50kDa kDa

Ubxn6 antibody - middle region - Additional Information**Gene ID** 66530**Alias Symbol** 1200008L11Rik, 2210415J11Rik, AU040909, Ubxd1, Ubxdc2**Other Names**

UBX domain-containing protein 6, UBX domain-containing protein 1, Ubxn6, Ubxd1, Ubxdc2

Format

Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.

Reconstitution & Storage

Add 50 ul of distilled water. Final anti-Ubxn6 antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.

Precautions

Ubxn6 antibody - middle region is for research use only and not for use in diagnostic or therapeutic procedures.

Ubxn6 antibody - middle region - Protein Information**Name** Ubxn6 {ECO:0000312|MGI:MGI:1913780}**Function**

May negatively regulate the ATPase activity of VCP, an ATP- driven segregase that associates with different cofactors to control a wide variety of cellular processes. As a cofactor of VCP, it may play a role in the transport of CAV1 to lysosomes for degradation. It may also play a role in endoplasmic reticulum-associated degradation (ERAD) of misfolded proteins. Together with VCP and other cofactors, it may play a role in macroautophagy, regulating for instance the clearance of damaged lysosomes.

Cellular Location

Cytoplasm {ECO:0000250|UniProtKB:Q9BZV1}. Cytoplasm, cytosol {ECO:0000250|UniProtKB:Q9BZV1}. Membrane {ECO:0000250|UniProtKB:Q9BZV1}; Peripheral membrane protein {ECO:0000250|UniProtKB:Q9BZV1}. Nucleus {ECO:0000250|UniProtKB:Q9BZV1}. Cytoplasm, cytoskeleton, microtubule organizing center, centrosome {ECO:0000250|UniProtKB:Q9BZV1}. Early endosome membrane {ECO:0000250|UniProtKB:Q9BZV1}; Peripheral membrane protein {ECO:0000250|UniProtKB:Q9BZV1}. Late endosome membrane {ECO:0000250|UniProtKB:Q9BZV1}; Peripheral membrane protein {ECO:0000250|UniProtKB:Q9BZV1}. Lysosome membrane {ECO:0000250|UniProtKB:Q9BZV1}; Peripheral membrane protein {ECO:0000250|UniProtKB:Q9BZV1}. Note=Localizes at the centrosome both in interphase and during mitosis. May be recruited to endosomal and lysosomal membranes as part of a ternary complex with CAV1 and VCP Recruited to damaged lysosomes decorated with K48-linked ubiquitin chains. {ECO:0000250|UniProtKB:Q9BZV1}

Tissue Location


Widely expressed (at protein level). Highest expression in brain (at protein level).

Ubxn6 antibody - middle region - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Ubxn6 antibody - middle region - Images



WB Suggested Anti-Ubxn6 Antibody Titration: 1.0 µg/ml

Positive Control: Mouse Heart

Ubxn6 antibody - middle region - References

- Carim-Todd L., et al. Biochim. Biophys. Acta 1517:298-301(2001).
Carninci P., et al. Science 309:1559-1563(2005).
Mural R.J., et al. Submitted (JUL-2005) to the EMBL/GenBank/DDBJ databases.
Church D.M., et al. PLoS Biol. 7:E1000112-E1000112(2009).
Madsen L., et al. Int. J. Biochem. Cell Biol. 40:2927-2942(2008).

