10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

USP38 antibody - middle region
Rabbit Polyclonal Antibody
Catalog # AI14382

Specification

USP38 antibody - middle region - Product Information

Application	**WB**
Primary Accession	Q8NB14
Other Accession	NM_032557, NP_115946
Reactivity	**Human, Mouse, Rat, Rabbit, Pig, Horse, Bovine, Guinea Pig, Dog**
Predicted	**Human, Mouse, Rabbit, Horse, Bovine, Dog**
Host	**Rabbit**
Clonality	**Polyclonal**
Calculated MW	**116kDa KDa**

USP38 antibody - middle region - Additional Information

Gene ID 84640

Alias Symbol	**FLJ35970, HP43.8KD, KIAA1891**

Other Names
Ubiquitin carboxyl-terminal hydrolase 38, 3.4.19.12, Deubiquitinating enzyme 38, HP43.8KD, Ubiquitin thioesterase 38, Ubiquitin-specific-processing protease 38, USP38, KIAA1891

Format
Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.

Reconstitution & Storage
Add 50 ul of distilled water. Final anti-USP38 antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.

Precautions
USP38 antibody - middle region is for research use only and not for use in diagnostic or therapeutic procedures.

USP38 antibody - middle region - Protein Information

Name USP38

Synonyms KIAA1891

Function
Deubiquitinating enzyme that plays a role in various cellular processes, including DNA repair, cell cycle regulation, and immune response (PubMed:22689415, PubMed:30497519, PubMed:31874856, PubMed:<a

href="http://www.uniprot.org/citations/35238669" target="_blank">35238669). Plays a role in the inhibition of type I interferon signaling by mediating the 'Lys-33' to 'Lys-48' ubiquitination transition of TBK1 leading to its degradation (PubMed:27692986). Cleaves the ubiquitin chain from the histone demethylase LSD1/KDM1A and prevents it from degradation by the 26S proteasome, thus maintaining LSD1 protein level in cells (PubMed:30497519). Plays a role in the DNA damage response by regulating the deacetylase activity of HDAC1 (PubMed:31874856). Mechanistically, removes the 'Lys-63'-linked ubiquitin chain promoting the deacetylase activity of HDAC1 in response to DNA damage (PubMed:31874856). Acts also as a specific deubiquitinase of histone deacetylase 3/HDAC3 and cleaves its 'Lys-63'-linked ubiquitin chains to lower its histone deacetylase activity (PubMed:32404892). Regulates MYC levels and cell proliferation via antagonizing ubiquitin E3 ligase FBXW7 thereby preventing MYC 'Lys-48'- linked ubiquitination and degradation (PubMed:34102342). Participates in antiviral response by removing both 'Lys-48'-linked and 'Lys-63'- linked polyubiquitination of Zika virus envelope protein E (PubMed:34696459). Constitutively associated with IL-33R/IL1RL1, deconjugates its 'Lys-27'-linked polyubiquitination resulting in its autophagic degradation (PubMed:35238669).

Cellular Location
Cytoplasm. Nucleus Note=In response to DNA damage, recruited to DNA damage sites in the nucleus.

Tissue Location
Highly expressed in skeletal muscle. Expressed in adrenal gland.


USP38 antibody - middle region - Protocols

Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- Cell Culture

USP38 antibody - middle region - Images

10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999



WB Suggested Anti-USP38 Antibody Titration: 0.2-1 µg/ml
ELISA Titer: 1:1562500
Positive Control: HepG2 cell lysate

USP38 antibody - middle region - References

Ota T.,et al.Nat. Genet. 36:40-45(2004).
Bechtel S.,et al.BMC Genomics 8:399-399(2007).
Hillier L.W.,et al.Nature 434:724-731(2005).
Mural R.J.,et al.Submitted (SEP-2005) to the EMBL/GenBank/DDBJ databases.
Nagase T.,et al.DNA Res. 8:179-187(2001).