

OR2D3 antibody - C-terminal region
Rabbit Polyclonal Antibody
Catalog # AI14445**Specification**

OR2D3 antibody - C-terminal region - Product Information

Application	WB
Primary Accession	Q8NGH3
Other Accession	NM_001004684 , NP_001004684
Reactivity	Human, Mouse, Rat, Pig, Horse, Bovine, Dog
Predicted	Human, Bovine
Host	Rabbit
Clonality	Polyclonal
Calculated MW	37kDa KDa

OR2D3 antibody - C-terminal region - Additional Information**Gene ID** 120775**Alias Symbol** OR11-89**Other Names**

Olfactory receptor 2D3, Olfactory receptor OR11-89, OR2D3

Format

Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.

Reconstitution & Storage

Add 50 ul of distilled water. Final anti-OR2D3 antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.

Precautions

OR2D3 antibody - C-terminal region is for research use only and not for use in diagnostic or therapeutic procedures.

OR2D3 antibody - C-terminal region - Protein Information**Name** OR2D3**Function**

Odorant receptor.

Cellular Location


Cell membrane; Multi-pass membrane protein.

OR2D3 antibody - C-terminal region - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

OR2D3 antibody - C-terminal region - Images



WB Suggested Anti-OR2D3 Antibody Titration: 1.0 µg/ml

Positive Control: 293T Whole Cell

OR2D3 antibody - C-terminal region - References

- Suwa M., et al. Submitted (JUL-2001) to the EMBL/GenBank/DDBJ databases.
Fuchs T., et al. Genomics 80:295-302(2002).
Malnic B., et al. Proc. Natl. Acad. Sci. U.S.A. 101:2584-2589(2004).
Malnic B., et al. Proc. Natl. Acad. Sci. U.S.A. 101:7205-7205(2004).