10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

Fyttd1 Antibody - middle region
Rabbit Polyclonal Antibody
Catalog # AI14520

Specification

Fyttd1 Antibody - middle region - Product Information

Application	**WB**
Primary Accession	Q7TQ84
Other Accession	NM_001047899, NP_001041364
Reactivity	**Human, Mouse, Rat, Rabbit, Pig, Horse, Bovine, Dog**
Predicted	**Human, Mouse, Rat, Rabbit, Pig, Horse, Bovine, Dog**
Host	**Rabbit**
Clonality	**Polyclonal**
Calculated MW	**20kDa KDa**

Fyttd1 Antibody - middle region - Additional Information

Alias Symbol	**RGD1306899**

Other Names
UAP56-interacting factor, Forty-two-three domain-containing protein 1, Protein 40-2-3, Fyttd1, Uif

Format
Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.

Reconstitution & Storage
Add 50 ul of distilled water. Final anti-Fyttd1 antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.

Precautions
Fyttd1 Antibody - middle region is for research use only and not for use in diagnostic or therapeutic procedures.

Fyttd1 Antibody - middle region - Protein Information

Name Fyttd1

Synonyms Uif

Function
Required for mRNA export from the nucleus to the cytoplasm. Acts as an adapter that uses the DDX39B/UAP56-NFX1 pathway to ensure efficient mRNA export and delivering to the nuclear pore. Associates with spliced and unspliced mRNAs simultaneously with ALYREF/THOC4 (By similarity).

Cellular Location
Nucleus, nucleoplasm. Nucleus speckle

10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

Fyttd1 Antibody - middle region - Protocols

Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- Cell Culture

Fyttd1 Antibody - middle region - Images



Host: Rabbit
Target Name: Fyttd1
Sample Tissue: Rat Thymus lysates
Antibody Dilution: 1.0µg/ml