

CD99L2 antibody - middle region
Rabbit Polyclonal Antibody
Catalog # AI14540**Specification**

CD99L2 antibody - middle region - Product Information

Application	WB
Primary Accession	Q8TCZ2
Other Accession	NM_134446 , NP_604395
Reactivity	Human, Mouse, Rat, Rabbit, Pig, Horse, Bovine, Guinea Pig, Dog
Predicted Host	Human, Mouse, Rabbit, Pig, Horse, Dog
Clonality	Rabbit
Calculated MW	Polyclonal 23kDa KDa

CD99L2 antibody - middle region - Additional Information**Gene ID** 83692**Alias Symbol** CD99B, DKFZp761H2024, MIC2L1**Other Names**

CD99 antigen-like protein 2, MIC2-like protein 1, CD99, CD99L2, MIC2L1

Format

Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.

Reconstitution & Storage

Add 50 ul of distilled water. Final anti-CD99L2 antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.

Precautions

CD99L2 antibody - middle region is for research use only and not for use in diagnostic or therapeutic procedures.

CD99L2 antibody - middle region - Protein Information**Name** CD99L2**Synonyms** MIC2L1**Function**

Plays a role in a late step of leukocyte extravasation helping cells to overcome the endothelial basement membrane. Acts at the same site as, but independently of, PECAM1 (By similarity). Homophilic adhesion molecule, but these interactions may not be required for cell aggregation (By similarity).

Cellular Location

Cell membrane; Single-pass type I membrane protein; Extracellular side. Cell junction. Secreted


Tissue Location

Detected in cerebrospinal fluid (at protein level) (PubMed:25326458). Expressed in many tissues, with low expression in thymus.

CD99L2 antibody - middle region - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

CD99L2 antibody - middle region - Images

WB Suggested Anti-CD99L2 Antibody Titration: 1.0 µg/ml
Positive Control: PANC1 Whole Cell