

FAM120C antibody - N-terminal region
Rabbit Polyclonal Antibody
Catalog # AI14560**Specification**

FAM120C antibody - N-terminal region - Product Information

Application	WB
Primary Accession	O9NX05
Other Accession	NM_017848 , NP_060318
Reactivity	Human, Mouse, Rat, Rabbit, Horse, Bovine, Dog
Predicted	Human, Mouse, Rat, Rabbit, Pig, Bovine, Dog
Host	Rabbit
Clonality	Polyclonal
Calculated MW	120kDa KDa

FAM120C antibody - N-terminal region - Additional Information**Gene ID** 54954**Alias Symbol** CXorf17, FLJ20506, FLJ43676, ORF34**Other Names**

Constitutive coactivator of PPAR-gamma-like protein 2, Protein FAM120C, Tumor antigen BJ-HCC-21, FAM120C, CXorf17

Format

Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.

Reconstitution & Storage

Add 50 ul of distilled water. Final anti-FAM120C antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.

Precautions

FAM120C antibody - N-terminal region is for research use only and not for use in diagnostic or therapeutic procedures.

FAM120C antibody - N-terminal region - Protein Information**Name** FAM120C**Synonyms** CXorf17**Tissue Location**


Expressed at low levels in a number of tissues.

FAM120C antibody - N-terminal region - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

FAM120C antibody - N-terminal region - Images



WB Suggested Anti-FAM120C Antibody Titration: 1.0 μ g/ml

Positive Control: Jurkat Whole Cell

FAM120C antibody - N-terminal region - References

- Holden S.,et al.Gene 318:149-161(2003).
Dong X.-Y.,et al.Submitted (JUN-2002) to the EMBL/GenBank/DDBJ databases.
Ota T.,et al.Nat. Genet. 36:40-45(2004).
Ross M.T.,et al.Nature 434:325-337(2005).