10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

EIF6 antibody - N-terminal region
Rabbit Polyclonal Antibody
Catalog # AI14664

Specification

EIF6 antibody - N-terminal region - Product Information

Application	**WB**
Primary Accession	P56537
Other Accession	NM_002212, NP_002203
Reactivity	**Human, Mouse, Rat, Rabbit, Sheep, Horse, Yeast, Bovine, Guinea Pig, Dog**
Predicted	**Human, Mouse, Chicken, Sheep, Guinea Pig, Dog**
Host	**Rabbit**
Clonality	**Polyclonal**
Calculated MW	**27kDa KDa**

EIF6 antibody - N-terminal region - Additional Information

Gene ID 3692

Alias Symbol	**2, CAB, EIF3A, ITGB4BP, b, b(2)gcn, gcn, p27BBP, eIF-6, p27(BBP)**

Other Names
Eukaryotic translation initiation factor 6 {ECO:0000255|HAMAP-Rule:MF_03132}, eIF-6 {ECO:0000255|HAMAP-Rule:MF_03132}, B(2)GCN homolog, B4 integrin interactor, CAB, p27(BBP), EIF6 {ECO:0000255|HAMAP-Rule:MF_03132}

Format
Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.

Reconstitution & Storage
Add 50 ul of distilled water. Final anti-EIF6 antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.

Precautions
EIF6 antibody - N-terminal region is for research use only and not for use in diagnostic or therapeutic procedures.

EIF6 antibody - N-terminal region - Protein Information

Name EIF6 {ECO:0000255|HAMAP-Rule:MF_03132, ECO:0000312|HGNC:HGNC:6159}

Function
Binds to the 60S ribosomal subunit and prevents its association with the 40S ribosomal subunit to form the 80S initiation complex in the cytoplasm (PubMed:10085284, PubMed:14654845, PubMed:<a

href="http://www.uniprot.org/citations/21536732" target="_blank">21536732, PubMed:32669547). Behaves as a stimulatory translation initiation factor downstream insulin/growth factors. Is also involved in ribosome biogenesis. Associates with pre-60S subunits in the nucleus and is involved in its nuclear export. Cytoplasmic release of TIF6 from 60S subunits and nuclear relocalization is promoted by a RACK1 (RACK1)- dependent protein kinase C activity (PubMed:10085284, PubMed:14654845, PubMed:21536732). In tissues responsive to insulin, controls fatty acid synthesis and glycolysis by exerting translational control of adipogenic transcription factors such as CEBPB, CEBPD and ATF4 that have G/C rich or uORF in their 5'UTR. Required for ROS-dependent megakaryocyte maturation and platelets formation, controls the expression of mitochondrial respiratory chain genes involved in reactive oxygen species (ROS) synthesis (By similarity). Involved in miRNA-mediated gene silencing by the RNA-induced silencing complex (RISC). Required for both miRNA-mediated translational repression and miRNA-mediated cleavage of complementary mRNAs by RISC (PubMed:17507929). Modulates cell cycle progression and global translation of pre-B cells, its activation seems to be rate-limiting in tumorigenesis and tumor growth (By similarity).

Cellular Location
Cytoplasm. Nucleus, nucleolus. Note=Shuttles between cytoplasm and nucleus/nucleolus


Tissue Location
Expressed at very high levels in colon carcinoma with lower levels in normal colon and ileum and lowest levels in kidney and muscle (at protein level).

EIF6 antibody - N-terminal region - Protocols

Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- Cell Culture


EIF6 antibody - N-terminal region - Images



WB Suggested Anti-EIF6 Antibody Titration: 1.0 µg/ml

10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

Positive Control: Hela Whole CellEIF6 is supported by BioGPS gene expression data to be expressed in HeLa



Host:Rabbit
Target Name:EIF6
Sample Tissue:Human 721_B
Antibody Dilution: 1.0µg/mlEIF6 is supported by BioGPS gene expression data to be expressed in 721_B

EIF6 antibody - N-terminal region - References

Si K.,et al.Proc. Natl. Acad. Sci. U.S.A. 94:14285-14290(1997).
Biffo S.,et al.J. Biol. Chem. 272:30314-30321(1997).
Donadini A.,et al.Gene 266:35-43(2001).
Mao M.,et al.Proc. Natl. Acad. Sci. U.S.A. 95:8175-8180(1998).
Ota T.,et al.Nat. Genet. 36:40-45(2004).