

RFC2 antibody - middle region
Rabbit Polyclonal Antibody
Catalog # AI14694**Specification**

RFC2 antibody - middle region - Product Information

Application	WB
Primary Accession	Q75MT5
Other Accession	NM_002914 , NP_002905
Reactivity	Human, Mouse, Rat, Rabbit, Horse, Bovine, Guinea Pig, Dog
Predicted	Human, Mouse, Rat, Rabbit, Chicken, Bovine, Dog
Host	Rabbit
Clonality	Polyclonal
Calculated MW	35kDa KDa

RFC2 antibody - middle region - Additional Information

Alias Symbol **A1, MGC3665, RFC40**

Format

Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.

Reconstitution & Storage

Add 50 ul of distilled water. Final anti-RFC2 antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.

Precautions


RFC2 antibody - middle region is for research use only and not for use in diagnostic or therapeutic procedures.

RFC2 antibody - middle region - Protein Information**RFC2 antibody - middle region - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

RFC2 antibody - middle region - Images



WB Suggested Anti-RFC2 Antibody Titration: 1.0 µg/ml

Positive Control: OVCAR-3 Whole Cell RFC2 is strongly supported by BioGPS gene expression data to be expressed in Human OVCAR3 cells