

NVL antibody - N-terminal region
Rabbit Polyclonal Antibody
Catalog # AI14696**Specification**

NVL antibody - N-terminal region - Product Information

Application	WB
Primary Accession	O15381-2
Other Accession	NM_206840 , NP_996671
Reactivity	Human, Rat, Pig, Horse, Bovine
Predicted	Human, Horse, Bovine
Host	Rabbit
Clonality	Polyclonal
Calculated MW	83kDa KDa

NVL antibody - N-terminal region - Additional Information**Format**

Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.

Reconstitution & Storage

Add 50 ul of distilled water. Final anti-NVL antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.


Precautions


NVL antibody - N-terminal region is for research use only and not for use in diagnostic or therapeutic procedures.

NVL antibody - N-terminal region - Protein Information**NVL antibody - N-terminal region - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

NVL antibody - N-terminal region - Images



WB Suggested Anti-NVL Antibody Titration: 1.0 µg/ml
Positive Control: THP-1 Whole Cell