

Mapre1 antibody - C-terminal region
Rabbit Polyclonal Antibody
Catalog # AI14714**Specification**

Mapre1 antibody - C-terminal region - Product Information

Application	WB
Primary Accession	Q61166
Other Accession	NM_007896 , NP_031922
Reactivity	Human, Mouse, Rat, Rabbit, Goat, Horse, Bovine, Guinea Pig, Dog
Predicted Host	Mouse, Pig, Dog
Clonality	Rabbit
Calculated MW	Polyclonal 30kDa KDa

Mapre1 antibody - C-terminal region - Additional Information**Gene ID** 13589**Alias Symbol** 5530600P05Rik, AI462499, AI504412, AW260097, BIM1p, D2Ert459e, Eb1**Other Names**

Microtubule-associated protein RP/EB family member 1, APC-binding protein EB1, End-binding protein 1, EB1, Mapre1

Format

Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.

Reconstitution & Storage

Add 50 ul of distilled water. Final anti-Mapre1 antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.

Precautions

Mapre1 antibody - C-terminal region is for research use only and not for use in diagnostic or therapeutic procedures.

Mapre1 antibody - C-terminal region - Protein Information**Name** Mapre1**Function**

Plus-end tracking protein (+TIP) that binds to the plus-end of microtubules and regulates the dynamics of the microtubule cytoskeleton. Promotes cytoplasmic microtubule nucleation and elongation. Involved in mitotic spindle positioning by stabilizing microtubules and promoting dynamic connection between astral microtubules and the cortex during mitotic chromosome segregation. Also acts as a regulator of minus-end microtubule organization: interacts with the complex formed by AKAP9 and PDE4DIP, leading to recruit CAMSAP2 to the Golgi apparatus,

thereby tethering non-centrosomal minus-end microtubules to the Golgi, an important step for polarized cell movement. Promotes elongation of CAMSAP2-decorated microtubule stretches on the minus-end of microtubules. Acts as a regulator of autophagosome transport via interaction with CAMSAP2 (By similarity). Functions downstream of Rho GTPases and DIAPH1 in stable microtubule formation (PubMed:15311282). May play a role in cell migration (PubMed:15311282).

Cellular Location

Cytoplasm, cytoskeleton. Cytoplasm, cytoskeleton, microtubule organizing center, centrosome {ECO:0000250|UniProtKB:Q15691} Cytoplasm, cytoskeleton, spindle {ECO:0000250|UniProtKB:Q15691} Cytoplasm, cytoskeleton, spindle pole {ECO:0000250|UniProtKB:Q15691} Note=Associated with the microtubule network at the growing distal tip of microtubules (PubMed:21357749). Also enriched at the centrosome (By similarity). {ECO:0000250|UniProtKB:Q15691, ECO:0000269|PubMed:21357749}

Tissue Location


Expressed within the midpiece of sperm tail (at protein level).

Mapre1 antibody - C-terminal region - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Mapre1 antibody - C-terminal region - Images



WB Suggested Anti-Mapre1 Antibody Titration: 1.0 µg/ml
Positive Control: Mouse Spleen

Mapre1 antibody - C-terminal region - References

Sparks A.B.,et al.Submitted (MAR-1996) to the EMBL/GenBank/DDBJ databases.
Bienvenut W.V.,et al.Submitted (FEB-2008) to UniProtKB.
Wen Y.,et al.Nat. Cell Biol. 6:820-830(2004).
Wu X.,et al.Cell 135:137-148(2008).

Fong K.W.,et al.Mol. Biol. Cell 20:3660-3670(2009).