

BAG4 antibody - C-terminal region
Rabbit Polyclonal Antibody
Catalog # AI14745**Specification**

BAG4 antibody - C-terminal region - Product Information

Application	WB
Primary Accession	B4E217
Other Accession	NM_001204878 , NP_001191807
Reactivity	Human, Rat, Rabbit, Pig, Horse, Bovine, Guinea Pig, Dog
Predicted Host	Human, Rat, Pig, Horse, Bovine, Dog
Clonality	Rabbit
Calculated MW	Polyclonal 46kDa KDa

BAG4 antibody - C-terminal region - Additional Information

Alias Symbol **BAG-4, SODD**

Format

Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.

Reconstitution & Storage

Add 50 ul of distilled water. Final anti-BAG4 antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.

Precautions


BAG4 antibody - C-terminal region is for research use only and not for use in diagnostic or therapeutic procedures.

BAG4 antibody - C-terminal region - Protein Information**BAG4 antibody - C-terminal region - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

BAG4 antibody - C-terminal region - Images



WB Suggested Anti-BAG4 Antibody Titration: 1.0 μ g/ml
Positive Control: THP-1 Whole Cell