

HOOK1 antibody - C-terminal region
Rabbit Polyclonal Antibody
Catalog # AI14770**Specification**

HOOK1 antibody - C-terminal region - Product Information

Application	WB
Primary Accession	O9UJC3
Other Accession	NM_015888 , NP_056972
Reactivity	Human, Mouse, Rat, Rabbit, Pig, Horse, Bovine, Guinea Pig, Dog
Predicted	Human, Mouse, Rat, Rabbit, Pig, Chicken, Horse, Bovine, Guinea Pig, Dog
Host	Rabbit
Clonality	Polyclonal
Calculated MW	80kDa KDa

HOOK1 antibody - C-terminal region - Additional Information**Gene ID** 51361

Alias Symbol	HK1, MGC10642
Other Names	
Protein Hook homolog 1, h-hook1, hHK1, HOOK1	

Format

Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.

Reconstitution & Storage

Add 50 ul of distilled water. Final anti-HOOK1 antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.

Precautions

HOOK1 antibody - C-terminal region is for research use only and not for use in diagnostic or therapeutic procedures.

HOOK1 antibody - C-terminal region - Protein Information**Name** HOOK1 ([HGNC:19884](#))**Function**

Component of the FTS/Hook/FHIP complex (FHF complex) (PubMed:18799622, PubMed:32073997). The FHF complex may function to promote vesicle trafficking and/or fusion via the homotypic vesicular protein sorting complex (the HOPS complex) (PubMed:18799622). FHF complex promotes the distribution of AP-4 complex to the perinuclear area of the cell (PubMed:18799622).

href="http://www.uniprot.org/citations/32073997" target="_blank">32073997). Required for spermatid differentiation. Probably involved in the positioning of the microtubules of the manchette and the flagellum in relation to the membrane skeleton (By similarity).

Cellular Location


Cytoplasm. Cytoplasm, cytoskeleton. Note=Localizes to punctate cytoplasmic foci which do not appear to overlap with early or late endosomes, the endoplasmic reticulum, multivesicular bodies (MVBs), lysosomes, or mitochondria (By similarity). Often found in close association with microtubules (By similarity). Does not associate with the Golgi complex. During spermiogenesis, it localizes to the manchette in spermatids from steps 8-10. It is also present between the microtubule manchette and the nucleus. During manchette elongation, it is preferentially localized to the nuclear ring of the manchette, whereas the strong localization to the manchette decreases. In more mature spermatids, while the manchette migrates posteriorly, it localizes to punctuates spots. At later stages of spermatid differentiation, the punctuate expression pattern is found at both the attachment site and the proximal end of the elongated manchette. In contrast, it is not present in mature spermatozoa (By similarity) {ECO:0000250|UniProtKB:Q8BIL5}

HOOK1 antibody - C-terminal region - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

HOOK1 antibody - C-terminal region - Images



WB Suggested Anti-HOOK1 Antibody Titration: 1.0 µg/ml
Positive Control: Fetal Heart

HOOK1 antibody - C-terminal region - References

Kraemer H., et al. Genetics 151:675-684(1999).
Ota T., et al. Nat. Genet. 36:40-45(2004).
Gregory S.G., et al. Nature 441:315-321(2006).
Mural R.J., et al. Submitted (SEP-2005) to the EMBL/GenBank/DDBJ databases.
Walenta J.H., et al. J. Cell Biol. 152:923-934(2001).