

MRPL40 antibody - middle region
Rabbit Polyclonal Antibody
Catalog # AI14830**Specification**

MRPL40 antibody - middle region - Product Information

Application	WB
Primary Accession	P15927
Other Accession	NM_002946 , NP_002937
Reactivity	Human, Mouse, Rat, Rabbit, Horse, Guinea Pig
Predicted Host	Human, Guinea Pig
Clonality	Rabbit
Calculated MW	Polyclonal 23kDa KDa

MRPL40 antibody - middle region - Additional Information**Gene ID** 6118

Alias Symbol **FLJ41774, MGC9400, MRP-L22, NLVCF, URIM**

Other Names

Replication protein A 32 kDa subunit, RP-A p32, Replication factor A protein 2, RF-A protein 2, Replication protein A 34 kDa subunit, RP-A p34, RPA2, REPA2, RPA32, RPA34

Format

Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.

Reconstitution & Storage

Add 50 ul of distilled water. Final anti-MRPL40 antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.

Precautions

MRPL40 antibody - middle region is for research use only and not for use in diagnostic or therapeutic procedures.

MRPL40 antibody - middle region - Protein Information

Name RPA2

Synonyms REPA2, RPA32, RPA34

Function

As part of the heterotrimeric replication protein A complex (RPA/RP-A), binds and stabilizes single-stranded DNA intermediates, that form during DNA replication or upon DNA stress. It prevents their reannealing and in parallel, recruits and activates different proteins and complexes involved in DNA metabolism. Thereby, it plays an essential role both in DNA replication and the

cellular response to DNA damage. In the cellular response to DNA damage, the RPA complex controls DNA repair and DNA damage checkpoint activation. Through recruitment of ATRIP activates the ATR kinase a master regulator of the DNA damage response. It is required for the recruitment of the DNA double-strand break repair factors RAD51 and RAD52 to chromatin in response to DNA damage. Also recruits to sites of DNA damage proteins like XPA and XPG that are involved in nucleotide excision repair and is required for this mechanism of DNA repair. Also plays a role in base excision repair (BER) probably through interaction with UNG. Also recruits SMARCAL1/HARP, which is involved in replication fork restart, to sites of DNA damage. May also play a role in telomere maintenance.

Cellular Location


Nucleus. Nucleus, PML body. Note=Redistributes to discrete nuclear foci upon DNA damage in an ATR-dependent manner

MRPL40 antibody - middle region - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

MRPL40 antibody - middle region - Images



WB Suggested Anti-MRPL40 Antibody Titration: 1.0 µg/ml
Positive Control: Fetal Liver

MRPL40 antibody - middle region - References

Erdile L.F.,et al.J. Biol. Chem. 265:3177-3182(1990).
Ebert L.,et al.Submitted (MAY-2004) to the EMBL/GenBank/DDBJ databases.
Gregory S.G.,et al.Nature 441:315-321(2006).
Din S.,et al.Genes Dev. 4:968-977(1990).
Dutta A.,et al.EMBO J. 11:2189-2199(1992).