

EDEM3 antibody - C-terminal region
Rabbit Polyclonal Antibody
Catalog # AI14834**Specification**

EDEM3 antibody - C-terminal region - Product Information

Application	WB
Primary Accession	O9P1W8-4
Other Accession	NM_001039508 , NP_001034597
Reactivity	Human, Mouse, Rat, Pig, Horse, Bovine, Dog
Predicted	Human, Mouse, Rat, Pig, Horse, Bovine, Dog
Host	Rabbit
Clonality	Polyclonal
Calculated MW	98kDa KDa

EDEM3 antibody - C-terminal region - Additional Information

Alias Symbol **C1orf22**

Format

Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.

Reconstitution & Storage

Add 50 ul of distilled water. Final anti-EDEM3 antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.

Precautions


EDEM3 antibody - C-terminal region is for research use only and not for use in diagnostic or therapeutic procedures.

EDEM3 antibody - C-terminal region - Protein Information**EDEM3 antibody - C-terminal region - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

EDEM3 antibody - C-terminal region - Images



WB Suggested Anti-EDEM3 Antibody Titration: 1.0 μ g/ml
Positive Control: Placenta