

TBRG4 antibody - C-terminal region
Rabbit Polyclonal Antibody
Catalog # AI14891**Specification**

TBRG4 antibody - C-terminal region - Product Information

Application	WB
Primary Accession	O969Z0-2
Other Accession	NM_030900 , NP_112162
Reactivity	Human, Rat, Pig, Bovine
Predicted	Human, Pig, Bovine
Host	Rabbit
Clonality	Polyclonal
Calculated MW	58kDa KDa

TBRG4 antibody - C-terminal region - Additional Information

Alias Symbol **CPR2, FASTKD4, KIAA0948**

Format

Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.

Reconstitution & Storage

Add 50 ul of distilled water. Final anti-TBRG4 antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.

Precautions


TBRG4 antibody - C-terminal region is for research use only and not for use in diagnostic or therapeutic procedures.

TBRG4 antibody - C-terminal region - Protein Information**TBRG4 antibody - C-terminal region - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

TBRG4 antibody - C-terminal region - Images



WB Suggested Anti-TBRG4 Antibody Titration: 1.0 µg/ml
Positive Control: MCF7 Whole Cell