

**RHD antibody - C-terminal region**  
**Rabbit Polyclonal Antibody**  
**Catalog # AI14916****Specification**

---

**RHD antibody - C-terminal region - Product Information**

|                   |                                                         |
|-------------------|---------------------------------------------------------|
| Application       | WB                                                      |
| Primary Accession | <a href="#">B2LR45</a>                                  |
| Other Accession   | <a href="#">NM_001127691</a> , <a href="#">ACB87205</a> |
| Reactivity        | Human, Bovine                                           |
| Predicted         | Human                                                   |
| Host              | Rabbit                                                  |
| Clonality         | Polyclonal                                              |
| Calculated MW     | 31kDa KDa                                               |

**RHD antibody - C-terminal region - Additional Information**

|              |                                                                                                                              |
|--------------|------------------------------------------------------------------------------------------------------------------------------|
| Alias Symbol | RP11-335G20.5, CD240D, DIIIc, MGC165007, RH, RH30, RHCED, RHDVA(TT), RHDel, RHPII, RHXIII, Rh4, RhDCw, RhII, RhK562-II, RhPI |
|--------------|------------------------------------------------------------------------------------------------------------------------------|

**Format**

Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.

**Reconstitution & Storage**

Add 50 ul of distilled water. Final anti-RHD antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.

**Precautions**


RHD antibody - C-terminal region is for research use only and not for use in diagnostic or therapeutic procedures.

**RHD antibody - C-terminal region - Protein Information****RHD antibody - C-terminal region - Protocols**

Provided below are standard protocols that you may find useful for product applications.


- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

## RHD antibody - C-terminal region - Images



WB Suggested Anti-RHD Antibody Titration: 1.0 µg/ml

Positive Control: RPMI-8226 Whole Cell. RHD is strongly supported by BioGPS gene expression data to be expressed in RPMI 8226




Host: Rabbit

Target Name: RHD

Sample Tissue: 293T

Antibody Dilution: 1.0 µg/ml RHD is strongly supported by BioGPS gene expression data to be expressed in Human HEK293T cells




Host: Rabbit


Target Name: RHD

Sample Tissue: Human Fetal Brain

Antibody Dilution: 1.0 µg/ml



Host:Rabbit  
Target Name:RHD  
Sample Tissue:Human Fetal Liver  
Antibody Dilution: 1.0µg/ml



Host:Rabbit  
Target Name:RHD  
Sample Tissue:Human Adult Placenta  
Antibody Dilution: 1.0µg/ml