10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

VILL antibody - C-terminal region
Rabbit Polyclonal Antibody
Catalog # AI14964

Specification

VILL antibody - C-terminal region - Product Information

Application	**WB**
Primary Accession	O15195
Other Accession	NM_015873, AAH00243
Reactivity	**Human, Pig, Horse, Bovine, Dog**
Predicted	**Human, Horse, Bovine, Dog**
Host	**Rabbit**
Clonality	**Polyclonal**
Calculated MW	**75kDa KDa**

VILL antibody - C-terminal region - Additional Information

Gene ID 50853

Other Names
Villin-like protein, VILL

Format
Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.

Reconstitution & Storage
Add 50 ul of distilled water. Final anti-VILL antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.

Precautions
VILL antibody - C-terminal region is for research use only and not for use in diagnostic or therapeutic procedures.

VILL antibody - C-terminal region - Protein Information

Name VILL

Function
Possible tumor suppressor.

Tissue Location
Ubiquitously expressed in 16 tissues examined.


VILL antibody - C-terminal region - Protocols

Provided below are standard protocols that you may find useful for product applications.

10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

- Western Blot
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- Cell Culture

VILL antibody - C-terminal region - Images



WB Suggested Anti-VILL Antibody Titration: 1.0 µg/ml
Positive Control: Fetal Liver

VILL antibody - C-terminal region - References

Muzny D.M.,et al.Nature 440:1194-1198(2006).
Mural R.J.,et al.Submitted (JUL-2005) to the EMBL/GenBank/DDBJ databases.
Ishikawa S.,et al.DNA Res. 4:35-43(1997).