

CNR2 antibody - C-terminal region
Rabbit Polyclonal Antibody
Catalog # AI14969**Specification**

CNR2 antibody - C-terminal region - Product Information

Application	WB
Primary Accession	P34972
Other Accession	NM_001841 , NP_001832
Reactivity	Human, Mouse, Rat, Horse
Predicted	Human, Mouse, Rat
Host	Rabbit
Clonality	Polyclonal
Calculated MW	40kDa KDa

CNR2 antibody - C-terminal region - Additional Information**Gene ID** 1269

Alias Symbol	CB2, CX5
Other Names	
Cannabinoid receptor 2, CB-2, CB2, hCB2, CX5, CNR2, CB2A, CB2B	

Format

Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.

Reconstitution & Storage

Add 50 ul of distilled water. Final anti-CNR2 antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.

Precautions

CNR2 antibody - C-terminal region is for research use only and not for use in diagnostic or therapeutic procedures.

CNR2 antibody - C-terminal region - Protein Information**Name** CNR2**Synonyms** CB2A, CB2B**Function**

Heterotrimeric G protein-coupled receptor for endocannabinoid 2-arachidonoylglycerol mediating inhibition of adenylate cyclase. May function in inflammatory response, nociceptive transmission and bone homeostasis.

Cellular Location

Cell membrane; Multi-pass membrane protein. Cell projection, dendrite. Perikaryon Note=Localizes to apical dendrite of pyramidal neurons.


Tissue Location

Preferentially expressed in cells of the immune system with higher expression in B-cells and NK cells (at protein level). Expressed in skin in suprabasal layers and hair follicles (at protein level). Highly expressed in tonsil and to a lower extent in spleen, peripheral blood mononuclear cells, and thymus. PubMed:14657172 could not detect expression in normal brain. Expressed in brain by perivascular microglial cells and dorsal root ganglion sensory neurons (at protein level). Two isoforms are produced by alternative promoter usage and differ only in the 5' UTR: isoform CB2A is observed predominantly in testis with some expression in brain, while isoform CB2B is predominant in spleen and leukocytes

CNR2 antibody - C-terminal region - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

CNR2 antibody - C-terminal region - Images

WB Suggested Anti-CNR2 Antibody Titration: 1.0 µg/ml

Positive Control: 721_B Whole Cell
CNR2 is supported by BioGPS gene expression data to be expressed in 721_B

CNR2 antibody - C-terminal region - References

- Munro S., et al. Nature 365:61-65(1993).
Liu Q.-R., et al. Genes Brain Behav. 8:519-530(2009).
Bruess M., et al. Submitted (FEB-2003) to the EMBL/GenBank/DDBJ databases.
Warren C.N., et al. Submitted (FEB-2003) to the EMBL/GenBank/DDBJ databases.
Saravanan T., et al. Submitted (NOV-2005) to the EMBL/GenBank/DDBJ databases.