

CSAD antibody - N-terminal region
Rabbit Polyclonal Antibody
Catalog # AI15083**Specification**

CSAD antibody - N-terminal region - Product Information

Application	WB
Primary Accession	O9Y602
Other Accession	NM_001244705 , AAD32543
Reactivity	Human, Mouse, Pig, Horse
Predicted	Human, Mouse, Pig, Horse
Host	Rabbit
Clonality	Polyclonal
Calculated MW	29kDa KDa

CSAD antibody - N-terminal region - Additional Information

Alias Symbol **CSD, PCAP**

Format

Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.

Reconstitution & Storage

Add 50 ul of distilled water. Final anti-CSAD antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.

Precautions


CSAD antibody - N-terminal region is for research use only and not for use in diagnostic or therapeutic procedures.

CSAD antibody - N-terminal region - Protein Information**CSAD antibody - N-terminal region - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

CSAD antibody - N-terminal region - Images



90 kDa
65 kDa
40 kDa
29 kDa
22 kDa

WB Suggested Anti-CSAD Antibody Titration: 1.0 µg/ml
Positive Control: 721_B Whole Cell