

**FKBP10 antibody - C-terminal region**  
**Rabbit Polyclonal Antibody**  
**Catalog # AI15186****Specification**

---

**FKBP10 antibody - C-terminal region - Product Information**

|                   |                                                                |
|-------------------|----------------------------------------------------------------|
| Application       | WB                                                             |
| Primary Accession | <a href="#">Q96AY3</a>                                         |
| Other Accession   | <a href="#">NM_021939</a> , <a href="#">NP_068758</a>          |
| Reactivity        | Human, Mouse, Rat, Rabbit, Horse, Bovine, Guinea Pig, Dog      |
| Predicted         | Human, Mouse, Rat, Rabbit, Pig, Horse, Bovine, Guinea Pig, Dog |
| Host              | Rabbit                                                         |
| Clonality         | Polyclonal                                                     |
| Calculated MW     | 65kDa kDa                                                      |

**FKBP10 antibody - C-terminal region - Additional Information****Gene ID** 60681**Alias Symbol** FKBP65, FLJ20683, FLJ22041, FLJ23833, PPIASE, hFKBP65, OI6, OI11**Other Names**

Peptidyl-prolyl cis-trans isomerase FKBP10, PPIase FKBP10, 5.2.1.8, 65 kDa FK506-binding protein, 65 kDa FKBP, FKBP-65, FK506-binding protein 10, FKBP-10, Immunophilin FKBP65, Rotamase, FKBP10, FKBP65

**Format**

Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.

**Reconstitution & Storage**

Add 50 ul of distilled water. Final anti-FKBP10 antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.

**Precautions**

FKBP10 antibody - C-terminal region is for research use only and not for use in diagnostic or therapeutic procedures.

**FKBP10 antibody - C-terminal region - Protein Information****Name** FKBP10**Synonyms** FKBP65**Function**

PPlases accelerate the folding of proteins during protein synthesis.


**Cellular Location**

Endoplasmic reticulum lumen {ECO:0000255|PROSITE- ProRule:PRU10138}

**FKBP10 antibody - C-terminal region - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

**FKBP10 antibody - C-terminal region - Images**

Host: Rabbit

Target Name: FKBP10

Antibody Dilution: 1.0µg/ml

Sample Tissue: PANC1 cell lysate

FKBP10 is supported by BioGPS gene expression data to be expressed in PANC1

**FKBP10 antibody - C-terminal region - References**

Rulten S.,et al.Submitted (JAN-2001) to the EMBL/GenBank/DDBJ databases.

Ota T.,et al.Nat. Genet. 36:40-45(2004).

Otsuki T.,et al.DNA Res. 12:117-126(2005).

Bechtel S.,et al.BMC Genomics 8:399-399(2007).

Zody M.C.,et al.Nature 440:1045-1049(2006).