

**CD300C Antibody - C-terminal region**  
**Rabbit Polyclonal Antibody**  
**Catalog # AI15203****Specification**

---

**CD300C Antibody - C-terminal region - Product Information**

|                   |                                                       |
|-------------------|-------------------------------------------------------|
| Application       | WB                                                    |
| Primary Accession | <a href="#">Q08708</a>                                |
| Other Accession   | <a href="#">NM_006678</a> , <a href="#">NP_006669</a> |
| Reactivity        | Human                                                 |
| Predicted         | Human                                                 |
| Host              | Rabbit                                                |
| Clonality         | Polyclonal                                            |
| Calculated MW     | 23kDa KDa                                             |

**CD300C Antibody - C-terminal region - Additional Information****Gene ID** 10871**Alias Symbol** CMRF-35A, CMRF35, CMRF35A, CMRF35A1, IGSF16, LIR**Other Names**

CMRF35-like molecule 6, CLM-6, CD300 antigen-like family member C, CMRF35-A1, CMRF-35, Immunoglobulin superfamily member 16, IgSF16, CD300c, CD300C, CMRF35, CMRF35A, CMRF35A1, IGSF16

**Format**

Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.

**Reconstitution & Storage**

Add 50 ul of distilled water. Final anti-CD300C antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.

**Precautions**

CD300C Antibody - C-terminal region is for research use only and not for use in diagnostic or therapeutic procedures.

**CD300C Antibody - C-terminal region - Protein Information****Name** CD300C**Synonyms** CMRF35, CMRF35A, CMRF35A1, IGSF16**Cellular Location**

Cell membrane; Single-pass type I membrane protein

**Tissue Location**

Present on the surface of monocytes, neutrophils, a proportion of peripheral blood T- and


B-lymphocytes and lymphocytic cell lines

### **CD300C Antibody - C-terminal region - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

### **CD300C Antibody - C-terminal region - Images**



WB Suggested Anti-CD300C Antibody Titration: 1.0 µg/ml  
Positive Control: 721\_B Whole Cell

### **CD300C Antibody - C-terminal region - References**

Jackson D.G., et al. Eur. J. Immunol. 22:1157-1163(1992).  
Clark G.J., et al. Tissue Antigens 57:415-423(2001).  
Speckman R.A., et al. Hum. Genet. 112:34-41(2003).