

CCDC26 Antibody - C-terminal region
Rabbit Polyclonal Antibody
Catalog # AI15250**Specification**

CCDC26 Antibody - C-terminal region - Product Information

Application	WB
Primary Accession	Q8TAB7
Other Accession	NM_145050 , NP_659487
Reactivity	Human
Predicted	Human
Host	Rabbit
Clonality	Polyclonal
Calculated MW	13kDa KDa

CCDC26 Antibody - C-terminal region - Additional Information

Alias Symbol **MGC27434, RAM**

Other Names

Putative coiled-coil domain-containing protein 26, Retinoic acid modulator, CCDC26, RAM

Format

Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.

Reconstitution & Storage

Add 50 ul of distilled water. Final anti-CCDC26 antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.


Precautions

CCDC26 Antibody - C-terminal region is for research use only and not for use in diagnostic or therapeutic procedures.

CCDC26 Antibody - C-terminal region - Protein Information**CCDC26 Antibody - C-terminal region - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

CCDC26 Antibody - C-terminal region - Images

WB Suggested Anti-CCDC26 Antibody Titration: 1.0 µg/ml
Positive Control: Fetal Liver

CCDC26 Antibody - C-terminal region - References

Nusbaum C., et al. Nature 439:331-335(2006).
Yin W., et al. Oncogene 25:3735-3744(2006).