

EXOC3L2 Antibody - C-terminal region
Rabbit Polyclonal Antibody
Catalog # AI15329**Specification**

EXOC3L2 Antibody - C-terminal region - Product Information

Application	WB
Primary Accession	Q2M3D2
Other Accession	NM_138568 , NP_612635
Reactivity	Human, Mouse, Rat, Bovine, Guinea Pig, Dog
Predicted	Human, Mouse, Rat, Pig, Bovine, Guinea Pig, Dog
Host	Rabbit
Clonality	Polyclonal
Calculated MW	46kDa KDa

EXOC3L2 Antibody - C-terminal region - Additional Information

Alias Symbol	FLJ36147, MGC16332, XTP7
Other Names	
Exocyst complex component 3-like protein 2, HBV X-transactivated gene 7 protein, HBV XAg-transactivated protein 7, EXOC3L2, XTP7	

Format

Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.

Reconstitution & Storage

Add 50 ul of distilled water. Final anti-EXOC3L2 antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.

Precautions

EXOC3L2 Antibody - C-terminal region is for research use only and not for use in diagnostic or therapeutic procedures.

EXOC3L2 Antibody - C-terminal region - Protein Information

Name EXOC3L2

Synonyms XTP7


EXOC3L2 Antibody - C-terminal region - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)

- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

EXOC3L2 Antibody - C-terminal region - Images



WB Suggested Anti-EXOC3L2 Antibody Titration: 1.0 µg/ml
Positive Control: Fetal Lung

EXOC3L2 Antibody - C-terminal region - References

Ota T., et al. Nat. Genet. 36:40-45(2004).
Mural R.J., et al. Submitted (JUL-2005) to the EMBL/GenBank/DDBJ databases.
Liu Y., et al. Submitted (MAR-2002) to the EMBL/GenBank/DDBJ databases.