10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

GPR31 Antibody - C-terminal region
Rabbit Polyclonal Antibody
Catalog # AI15364

Specification

GPR31 Antibody - C-terminal region - Product Information

Application	**WB**
Primary Accession	O00270
Other Accession	NP_005290
Reactivity	**Human**
Host	**Rabbit**
Clonality	**Polyclonal**
Calculated MW	**35kDa KDa**

GPR31 Antibody - C-terminal region - Additional Information

Gene ID 2853

Alias Symbol	**GPR31,**

Other Names
12-(S)-hydroxy-5, 8, 10, 14-eicosatetraenoic acid receptor, 12-(S)-HETE receptor, 12-HETER, G-protein coupled receptor 31, GPR31

Format
Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.

Reconstitution & Storage
Add 50 &mu, l of distilled water. Final Anti-GPR31 antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at -20°C. Avoid repeat freeze-thaw cycles.

Precautions
GPR31 Antibody - C-terminal region is for research use only and not for use in diagnostic or therapeutic procedures.

GPR31 Antibody - C-terminal region - Protein Information

Name GPR31

Function
High-affinity receptor for 12-(S)-hydroxy-5,8,10,14- eicosatetraenoic acid (12-S-HETE), with much lower affinities for other HETE isomers (PubMed:21712392, PubMed:29227475). 12-S-HETE is a eicosanoid, a 12-lipoxygenase (ALOX12) metabolite of arachidonic acid, involved in many physiologic and pathologic processes (PubMed:26965684, PubMed:28619714, PubMed:29227475).

12-S-HETE-binding leads to activation of ERK1/2 (MAPK3/MAPK1), MEK, and NF-kappa-B pathways leading to cell growth (PubMed:21712392, PubMed:29227475). Plays a crucial role for proliferation, survival and macropinocytosis of KRAS- dependent cancer cells by mediating the translocation of KRAS from the endoplasmic reticulum to the plasma membrane (PM) and its association with the PM (PubMed:28619714). Contributes to enhanced immune responses by inducing dendrite protrusion of small intestinal CX3CR1(+) phagocytes for the uptake of luminal antigens (By similarity). Acts also as a key receptor for 12-(S)-HETE-mediated liver ischemia reperfusion injury (PubMed:29227475).


Cellular Location
Cell membrane; Multi-pass membrane protein

GPR31 Antibody - C-terminal region - Protocols

Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- Cell Culture

GPR31 Antibody - C-terminal region - Images



Host: Rabbit
Target Name: GPR31
Sample Tissue: HepG2 Whole cell lysate
s
Antibody Dilution: 1.0µg/ml

GPR31 Antibody - C-terminal region - References

Zingoni A.,et al.Genomics 42:519-523(1997).
Kaighin V.A.,et al.Submitted (DEC-2007) to the EMBL/GenBank/DDBJ databases.
Mungall A.J.,et al.Nature 425:805-811(2003).
Mural R.J.,et al.Submitted (SEP-2005) to the EMBL/GenBank/DDBJ databases.
Guo Y.,et al.J. Biol. Chem. 286:33832-33840(2011).

10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999