

PAPLN Antibody - C-terminal region
Rabbit Polyclonal Antibody
Catalog # AI15373**Specification**

PAPLN Antibody - C-terminal region - Product Information

Application	WB
Primary Accession	O95428
Other Accession	NM_173462 , NP_775733
Reactivity	Human, Rabbit, Pig, Horse, Dog
Predicted	Human, Rabbit, Pig, Horse, Dog
Host	Rabbit
Clonality	Polyclonal
Calculated MW	138kDa KDa

PAPLN Antibody - C-terminal region - Additional Information**Gene ID** 89932

Alias Symbol	DKFZp434F053, MGC50452
Other Names	
Papilin, PAPLN	

Format

Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.

Reconstitution & Storage

Add 50 ul of distilled water. Final anti-PAPLN antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.

Precautions

PAPLN Antibody - C-terminal region is for research use only and not for use in diagnostic or therapeutic procedures.


PAPLN Antibody - C-terminal region - Protein Information**Name** PAPLN**Cellular Location**
Secreted.**PAPLN Antibody - C-terminal region - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)

- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

PAPLN Antibody - C-terminal region - Images



WB Suggested Anti-PAPLN Antibody Titration: 1.0 µg/ml
Positive Control: HepG2 Whole Cell

PAPLN Antibody - C-terminal region - References

Clark H.F., et al. Genome Res. 13:2265-2270(2003).
Ota T., et al. Nat. Genet. 36:40-45(2004).
Heilig R., et al. Nature 421:601-607(2003).
Bechtel S., et al. BMC Genomics 8:399-399(2007).
Jikuya H., et al. Submitted (SEP-2003) to the EMBL/GenBank/DDBJ databases.