

CRYGN Antibody - C-terminal region
Rabbit Polyclonal Antibody
Catalog # AI15426**Specification**

CRYGN Antibody - C-terminal region - Product Information

Application	WB
Primary Accession	Q8WXF5
Other Accession	NM_144727 , NP_653328
Reactivity	Human
Predicted	Human
Host	Rabbit
Clonality	Polyclonal
Calculated MW	20kDa KDa

CRYGN Antibody - C-terminal region - Additional Information**Gene ID** 155051**Alias Symbol** **MGC119042, MGC119043, MGC119044, MGC119045****Other Names**
Gamma-crystallin N, Gamma-N-crystallin, CRYGN**Format**

Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.

Reconstitution & Storage

Add 50 ul of distilled water. Final anti-CRYGN antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.

Precautions

CRYGN Antibody - C-terminal region is for research use only and not for use in diagnostic or therapeutic procedures.

CRYGN Antibody - C-terminal region - Protein Information**Name** CRYGN ([HGNC:20458](#))**Tissue Location**


Not specifically expressed in eye.

CRYGN Antibody - C-terminal region - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

CRYGN Antibody - C-terminal region - Images



WB Suggested Anti-CRYGN Antibody Titration: 1.0 µg/ml

Positive Control: HeLa Whole cell

CRYGN Antibody - C-terminal region - References

Wistow G., et al. FEBS J. 272:2276-2291(2005).

Mural R.J., et al. Submitted (SEP-2005) to the EMBL/GenBank/DDBJ databases.