

BTN3A1 Antibody - C-terminal region
Rabbit Polyclonal Antibody
Catalog # AI15476**Specification**

BTN3A1 Antibody - C-terminal region - Product Information

Application	WB
Primary Accession	O00481-3
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal
Calculated MW	41kDa KDa

BTN3A1 Antibody - C-terminal region - Additional Information

Alias Symbol **BTN3A1, BTF5,**
Format

Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.

Reconstitution & Storage

Add 50 µl of distilled water. Final Anti-BTN3A1 antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at -20°C. Avoid repeat freeze-thaw cycles.

Precautions

BTN3A1 Antibody - C-terminal region is for research use only and not for use in diagnostic or therapeutic procedures.

BTN3A1 Antibody - C-terminal region - Protein Information**BTN3A1 Antibody - C-terminal region - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

BTN3A1 Antibody - C-terminal region - Images

90 kDa__
65 kDa__
40 kDa__
29 kDa__
22 kDa__



Host: Rabbit
Target Name: BTN3A1
Sample Tissue: HepG2 Cell Lysate
Antibody Dilution: 1.0µg/ml

Host: Rabbit
Target Name: BTN3A1
Sample Tissue: HepG2 Whole cell lysate
s
Antibody Dilution: 1.0µg/ml