

F2RL3 Antibody - C-terminal region
Rabbit Polyclonal Antibody
Catalog # AI15491**Specification**

F2RL3 Antibody - C-terminal region - Product Information

Application	WB
Primary Accession	O96RIO
Other Accession	NM_003950 , NP_003941
Reactivity	Human
Predicted	Human
Host	Rabbit
Clonality	Polyclonal
Calculated MW	42kDa KDa

F2RL3 Antibody - C-terminal region - Additional Information**Gene ID** 9002**Alias Symbol** **PAR4****Other Names**

Proteinase-activated receptor 4, PAR-4, Coagulation factor II receptor-like 3, Thrombin receptor-like 3, F2RL3, PAR4

Format

Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.

Reconstitution & Storage

Add 50 ul of distilled water. Final anti-F2RL3 antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.

Precautions

F2RL3 Antibody - C-terminal region is for research use only and not for use in diagnostic or therapeutic procedures.

F2RL3 Antibody - C-terminal region - Protein Information**Name** F2RL3**Synonyms** PAR4**Function**

Receptor for activated thrombin or trypsin coupled to G proteins that stimulate phosphoinositide hydrolysis (PubMed:10079109). May play a role in platelets activation (PubMed:10079109).

Cellular Location

Cell membrane; Multi-pass membrane protein.

Tissue Location


Widely expressed, with highest levels in lung, pancreas, thyroid, testis and small intestine. Not expressed in brain, kidney, spinal cord and peripheral blood leukocytes. Also detected in platelets

F2RL3 Antibody - C-terminal region - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

F2RL3 Antibody - C-terminal region - Images



Host: Rabbit

Target Name: F2RL3

Sample Tissue: THP-1 Whole cell lysate

s

Antibody Dilution: 1.0µg/ml

F2RL3 Antibody - C-terminal region - References

Kahn M.L.,et al.J. Biol. Chem. 273:23290-23296(1998).

Kahn M.L.,et al.Nature 394:690-694(1998).

Xu W.-F.,et al.Proc. Natl. Acad. Sci. U.S.A. 95:6642-6646(1998).

King M.M.,et al.Submitted (OCT-2003) to the EMBL/GenBank/DDBJ databases.

Kahn M.L.,et al.J. Clin. Invest. 103:879-887(1999).