

TIGD7 Antibody - C-terminal region
Rabbit Polyclonal Antibody
Catalog # AI15540**Specification**

TIGD7 Antibody - C-terminal region - Product Information

Application	WB
Primary Accession	Q6NT04
Other Accession	NM_033208 , NP_149985
Reactivity	Human
Predicted	Human
Host	Rabbit
Clonality	Polyclonal
Calculated MW	60kDa KDa

TIGD7 Antibody - C-terminal region - Additional Information**Gene ID** 91151**Alias Symbol** **Sancho****Other Names**

Tigger transposable element-derived protein 7, TIGD7

Format

Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.

Reconstitution & Storage

Add 50 ul of distilled water. Final anti-TIGD7 antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.

Precautions

TIGD7 Antibody - C-terminal region is for research use only and not for use in diagnostic or therapeutic procedures.

TIGD7 Antibody - C-terminal region - Protein Information**Name** TIGD7**Cellular Location**

Nucleus.

Tissue Location


Expressed in all tissues tested. Higher expression in testis and ovary.

TIGD7 Antibody - C-terminal region - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

TIGD7 Antibody - C-terminal region - Images



Host: Rabbit

Target Name: TIGD7

Sample Tissue: Hela Whole cell lysate

S

Antibody Dilution: 1.0µg/ml

TIGD7 Antibody - C-terminal region - References

Dou T., et al. Biochem. Genet. 42:279-285(2004).