

CCDC86 Antibody - N-terminal region
Rabbit Polyclonal Antibody
Catalog # AI15584**Specification**

CCDC86 Antibody - N-terminal region - Product Information

Application	WB
Primary Accession	Q9H6F5
Other Accession	NM_024098 , NP_077003
Reactivity	Human, Horse, Bovine
Predicted	Human, Horse, Bovine
Host	Rabbit
Clonality	Polyclonal
Calculated MW	40kDa KDa

CCDC86 Antibody - N-terminal region - Additional Information**Gene ID** 79080**Other Names**

Coiled-coil domain-containing protein 86, Cytokine-induced protein with coiled-coil domain, CCDC86, CYCLON

Format

Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.

Reconstitution & Storage

Add 50 ul of distilled water. Final anti-CCDC86 antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.

Precautions

CCDC86 Antibody - N-terminal region is for research use only and not for use in diagnostic or therapeutic procedures.

CCDC86 Antibody - N-terminal region - Protein Information**Name** CCDC86**Synonyms** CYCLON**Cellular Location**


Nucleus.

CCDC86 Antibody - N-terminal region - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

CCDC86 Antibody - N-terminal region - Images



Host: Rabbit

Target Name: CCDC86

Sample Tissue: MCF7 Whole cell lysate

s

Antibody Dilution: 1.0µg/mlCCDC86 is supported by BioGPS gene expression data to be expressed in MCF7

CCDC86 Antibody - N-terminal region - References

Hoshino A.,et al.FEBS Lett. 581:975-980(2007).

Ota T.,et al.Nat. Genet. 36:40-45(2004).

Taylor T.D.,et al.Nature 440:497-500(2006).

Ahn J.,et al.J. Biochem. Mol. Biol. 37:741-748(2004).

Olsen J.V.,et al.Cell 127:635-648(2006).