10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

DCAF8 Antibody - C-terminal region
Rabbit Polyclonal Antibody
Catalog # AI15801

Specification

DCAF8 Antibody - C-terminal region - Product Information

Application	**WB**
Primary Accession	Q5TAQ9
Other Accession	NM_015726, NP_056541
Reactivity	**Human, Mouse, Rat, Rabbit, Pig, Horse, Bovine, Guinea Pig, Dog**
Predicted	**Human, Mouse, Rat, Rabbit, Pig, Horse, Bovine, Guinea Pig, Dog**
Host	**Rabbit**
Clonality	**Polyclonal**
Calculated MW	**67kDa KDa**

DCAF8 Antibody - C-terminal region - Additional Information

Gene ID 50717

Alias Symbol	**H326, WDR42A**

Other Names
DDB1- and CUL4-associated factor 8, WD repeat-containing protein 42A, DCAF8, H326, WDR42A

Format
Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.

Reconstitution & Storage
Add 50 ul of distilled water. Final anti-DCAF8 antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.

Precautions
DCAF8 Antibody - C-terminal region is for research use only and not for use in diagnostic or therapeutic procedures.

DCAF8 Antibody - C-terminal region - Protein Information

Name DCAF8

Synonyms H326, WDR42A

Function
May function as a substrate receptor for CUL4-DDB1 E3 ubiquitin-protein ligase complex.

Cellular Location
Nucleus. Cytoplasm. Note=It shuttles between the nucleus and the cytoplasm. Nuclear import is mediated by KPNA1 and KPNB1 under the regulation of nuclear GTPase RAN. Nuclear export to the


cytoplasm is XPO1 dependent.

DCAF8 Antibody - C-terminal region - Protocols

Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- Cell Culture

DCAF8 Antibody - C-terminal region - Images



Host: Rabbit
Target Name: DCAF8
Sample Tissue: 721_B Whole cell lysate
s
Antibody Dilution: 1.0µg/mlDCAF8 is strongly supported by BioGPS gene expression data to be expressed in Human 721_B cells

DCAF8 Antibody - C-terminal region - Background

May function as a substrate receptor for CUL4-DDB1 E3 ubiquitin-protein ligase complex.

DCAF8 Antibody - C-terminal region - References

Bergsagel P.L.,et al.Submitted (FEB-1994) to the EMBL/GenBank/DDBJ databases.
Totoki Y.,et al.Submitted (APR-2005) to the EMBL/GenBank/DDBJ databases.
Bechtel S.,et al.BMC Genomics 8:399-399(2007).
Gregory S.G.,et al.Nature 441:315-321(2006).
Mural R.J.,et al.Submitted (SEP-2005) to the EMBL/GenBank/DDBJ databases.