

OR2M2 Antibody - C-terminal region
Rabbit Polyclonal Antibody
Catalog # AI15907**Specification**

OR2M2 Antibody - C-terminal region - Product Information

Application	WB
Primary Accession	Q96R28
Other Accession	NM_001004688 , NP_001004688
Reactivity	Human, Pig, Horse, Bovine, Guinea Pig, Dog
Predicted	Human, Pig, Horse, Bovine, Guinea Pig, Dog
Host	Rabbit
Clonality	Polyclonal
Calculated MW	38kDa KDa

OR2M2 Antibody - C-terminal region - Additional Information**Gene ID** 391194

Alias Symbol	OR2M2Q, OST423
Other Names	
Olfactory receptor 2M2, OST423, OR2M2	

Format

Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.

Reconstitution & Storage

Add 50 ul of distilled water. Final anti-OR2M2 antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.

Precautions

OR2M2 Antibody - C-terminal region is for research use only and not for use in diagnostic or therapeutic procedures.

OR2M2 Antibody - C-terminal region - Protein Information**Name** OR2M2**Function**

Odorant receptor.

Cellular Location


Cell membrane; Multi-pass membrane protein.

OR2M2 Antibody - C-terminal region - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

OR2M2 Antibody - C-terminal region - Images



Host: Rabbit
Target Name: OR2M2
Sample Tissue: Jurkat Whole cell lysate
s
Antibody Dilution: 1.0µg/ml

OR2M2 Antibody - C-terminal region - Background

Odorant receptor.

OR2M2 Antibody - C-terminal region - References

Gregory S.G.,et al.Nature 441:315-321(2006).
Fuchs T.,et al.Genomics 80:295-302(2002).