

**CFI Antibody - C-terminal region**  
**Rabbit Polyclonal Antibody**  
**Catalog # AI15961****Specification**

---

**CFI Antibody - C-terminal region - Product Information**

Application	WB
Primary Accession	<a href="#">P05156</a>
Other Accession	<a href="#">NM_000204</a> , <a href="#">NP_000195</a>
Reactivity	Human, Mouse, Rat, Rabbit, Pig, Horse, Bovine, Guinea Pig, Dog
Predicted	Human, Mouse, Rat, Rabbit, Pig, Horse, Bovine, Guinea Pig, Dog
Host	Rabbit
Clonality	Polyclonal
Calculated MW	64kDa kDa

**CFI Antibody - C-terminal region - Additional Information****Gene ID** 3426**Alias Symbol** CFI, IF,**Other Names**

Complement factor I, 3.4.21.45, C3B/C4B inactivator, Complement factor I heavy chain, Complement factor I light chain, CFI, IF

**Format**

Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.

**Reconstitution & Storage**

Add 50 µl of distilled water. Final Anti-CFI antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at -20°C. Avoid repeat freeze-thaw cycles.

**Precautions**

CFI Antibody - C-terminal region is for research use only and not for use in diagnostic or therapeutic procedures.

**CFI Antibody - C-terminal region - Protein Information****Name** CFI**Synonyms** IF**Function**

Trypsin-like serine protease that plays an essential role in regulating the immune response by controlling all complement pathways. Inhibits these pathways by cleaving three peptide bonds in the alpha-chain of C3b and two bonds in the alpha-chain of C4b thereby inactivating these proteins (PubMed: &lt;a href="http://www.uniprot.org/citations/7360115"

target="\_blank">7360115</a>, PubMed:<a href="http://www.uniprot.org/citations/17320177" target="\_blank">17320177</a>). Essential cofactors for these reactions include factor H and C4BP in the fluid phase and membrane cofactor protein/CD46 and CR1 on cell surfaces (PubMed:<a href="http://www.uniprot.org/citations/2141838" target="\_blank">2141838</a>, PubMed:<a href="http://www.uniprot.org/citations/9605165" target="\_blank">9605165</a>, PubMed:<a href="http://www.uniprot.org/citations/12055245" target="\_blank">12055245</a>). The presence of these cofactors on healthy cells allows degradation of deposited C3b by CFI in order to prevent undesired complement activation, while in apoptotic cells or microbes, the absence of such cofactors leads to C3b-mediated complement activation and subsequent opsonization (PubMed:<a href="http://www.uniprot.org/citations/28671664" target="\_blank">28671664</a>).

### Cellular Location

Secreted, extracellular space. Secreted

### Tissue Location


Expressed in the liver by hepatocytes (PubMed:6327681). Also present in other cells such as monocytes, fibroblasts or keratinocytes (PubMed:6444659, PubMed:17320177)

## CFI Antibody - C-terminal region - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

## CFI Antibody - C-terminal region - Images



Host: Rabbit  
Target Name: CFI  
Sample Tissue: Jurkat Whole Cell lysates  
Antibody Dilution: 1.0µg/ml

## CFI Antibody - C-terminal region - Background

Responsible for cleaving the alpha-chains of C4b and C3b in the presence of the cofactors C4-binding protein and factor H respectively.

**CFI Antibody - C-terminal region - References**

Catterall C.F.,et al.Biochem. J. 242:849-856(1987).  
Goldberger G.,et al.J. Biol. Chem. 262:10065-10071(1987).  
Hillier L.W.,et al.Nature 434:724-731(2005).  
Minta J.O.,et al.Gene 208:17-24(1998).  
Ullman C.G.,et al.FEBS Lett. 371:199-203(1995).