10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

TSPYL5 Antibody - N-terminal region
Rabbit Polyclonal Antibody
Catalog # AI16072

Specification

TSPYL5 Antibody - N-terminal region - Product Information

Application	**WB**
Primary Accession	Q86VY4
Other Accession	NP_277047
Reactivity	**Human**
Host	**Rabbit**
Clonality	**Polyclonal**
Calculated MW	**45kDa KDa**

TSPYL5 Antibody - N-terminal region - Additional Information

Gene ID 85453

Alias Symbol **TSPYL5, KIAA1750,**
Other Names
Testis-specific Y-encoded-like protein 5, TSPY-like protein 5, TSPYL5, KIAA1750

Format
Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.

Reconstitution & Storage
Add 50 &mu, l of distilled water. Final Anti-TSPYL5 antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at -20°C. Avoid repeat freeze-thaw cycles.

Precautions
TSPYL5 Antibody - N-terminal region is for research use only and not for use in diagnostic or therapeutic procedures.

TSPYL5 Antibody - N-terminal region - Protein Information

Name TSPYL5

Synonyms KIAA1750

Function
Involved in modulation of cell growth and cellular response to gamma radiation probably via regulation of the Akt signaling pathway. Involved in regulation of p53/TP53. Suppresses p53/TP53 protein levels and promotes its ubiquitination; the function is dependent on USP7 and independent on MDM2. Proposed to displace p53/TP53 from interaction with USP7.


TSPYL5 Antibody - N-terminal region - Protocols

10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- Cell Culture

TSPYL5 Antibody - N-terminal region - Images



Host: Rabbit
Target Name: TSPYL5
Sample Tissue: Jurkat Whole cell lysate
s
Antibody Dilution: 1.0µg/ml

TSPYL5 Antibody - N-terminal region - Background

Involved in modulation of cell growth and cellular response to gamma radiation probably via regulation of the Akt signaling pathway. Involved in regulation of p53/TP53. Suppresses p53/TP53 protein levels and promotes its ubiquitination; the function is dependent on USP7 and independent on MDM2. Proposed to displace p53/TP53 from interaction with USP7.

TSPYL5 Antibody - N-terminal region - References

Nagase T.,et al.DNA Res. 7:347-355(2000).
Ota T.,et al.Nat. Genet. 36:40-45(2004).
Dephoure N.,et al.Proc. Natl. Acad. Sci. U.S.A. 105:10762-10767(2008).
Kim E.J.,et al.Biochem. Biophys. Res. Commun. 392:448-453(2010).
Burkard T.R.,et al.BMC Syst. Biol. 5:17-17(2011).