

MYPOP antibody - N-terminal region
Rabbit Polyclonal Antibody
Catalog # AI16232**Specification**

MYPOP antibody - N-terminal region - Product Information

| | |
|-------------------|---|
| Application | WB |
| Primary Accession | Q86VE0 |
| Other Accession | NM_001012643 , NP_001012661 |
| Reactivity | Human, Mouse, Rat, Pig, Bovine, Guinea Pig, Dog |
| Predicted | Human, Mouse, Rat, Pig, Bovine, Guinea Pig, Dog |
| Host | Rabbit |
| Clonality | Polyclonal |
| Calculated MW | 42kDa KDa |

MYPOP antibody - N-terminal region - Additional Information**Gene ID** 339344**Alias Symbol** **P42pop****Other Names**

Myb-related transcription factor, partner of profilin, Myb-related protein p42POP, Partner of profilin, MYPOP, P42POP

Format

Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.

Reconstitution & Storage

Add 50 ul of distilled water. Final anti-MYPOP antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.

Precautions

MYPOP antibody - N-terminal region is for research use only and not for use in diagnostic or therapeutic procedures.

MYPOP antibody - N-terminal region - Protein Information**Name** MYPOP**Synonyms** P42POP**Function**

Transcriptional repressor; DNA-binding protein that specifically recognizes the core sequence 5'-YAAC[GT]G-3'. Dimerization with PFN1 reduces its DNA-binding capacity (By similarity).

Cellular Location


Nucleus.

MYPOP antibody - N-terminal region - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

MYPOP antibody - N-terminal region - Images



WB Suggested Anti-MYPOP Antibody Titration: 0.2-1 μ g/ml

ELISA Titer: 1:62500

Positive Control: COLO205 cell lysate

MYPOP antibody - N-terminal region - Background

Transcriptional repressor; DNA-binding protein that specifically recognizes the core sequence 5'-YAAC[GT]G-3'. Dimerization with PFN1 reduces its DNA-binding capacity (By similarity).