

GTF2B antibody - N-terminal region
Rabbit Polyclonal Antibody
Catalog # AI16268**Specification**

GTF2B antibody - N-terminal region - Product Information

Application	WB
Primary Accession	Q00403
Other Accession	NM_001514 , NP_001505
Reactivity	Human, Mouse, Rabbit, Pig, Horse, Guinea Pig, Dog
Predicted	Human, Mouse, Rabbit, Pig, Chicken, Horse, Guinea Pig, Dog
Host	Rabbit
Clonality	Polyclonal
Calculated MW	35kDa KDa

GTF2B antibody - N-terminal region - Additional Information**Gene ID** 2959

Alias Symbol	TF2B, TFIIB
Other Names	
Transcription initiation factor IIB, General transcription factor TFIIB, S300-II, GTF2B, TF2B, TFIIB	

Format

Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.

Reconstitution & Storage

Add 50 ul of distilled water. Final anti-GTF2B antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.

Precautions

GTF2B antibody - N-terminal region is for research use only and not for use in diagnostic or therapeutic procedures.

GTF2B antibody - N-terminal region - Protein Information**Name** GTF2B**Synonyms** TF2B, TFIIB**Function**

General transcription factor that plays a role in transcription initiation by RNA polymerase II (Pol II). Involved in the pre-initiation complex (PIC) formation and Pol II recruitment at promoter DNA (PubMed: [1876184](http://www.uniprot.org/citations/1876184), PubMed: [1946368](http://www.uniprot.org/citations/1946368), PubMed: [1517211](http://www.uniprot.org/citations/1517211)).

PubMed:3818643, PubMed:3029109, PubMed:8413225, PubMed:8515820, PubMed:8516311, PubMed:8516312, PubMed:7601352, PubMed:9420329, PubMed:12931194, PubMed:27193682). Together with the TATA box-bound TBP forms the core initiation complex and provides a bridge between TBP and the Pol II-TFIIF complex (PubMed:8504927, PubMed:8413225, PubMed:8515820, PubMed:8516311, PubMed:8516312). Released from the PIC early following the onset of transcription during the initiation and elongation transition and reassociates with TBP during the next transcription cycle (PubMed:7601352). Associates with chromatin to core promoter-specific regions (PubMed:12931194, PubMed:24441171). Binds to two distinct DNA core promoter consensus sequence elements in a TBP- independent manner; these IIB-recognition elements (BREs) are localized immediately upstream (BREu), 5'-[GC][GC][GA]CGCC-3', and downstream (BREd), 5'-[GA]T[TGA][TG][GT][TG][TG]-3', of the TATA box element (PubMed:9420329, PubMed:16230532, PubMed:7675079, PubMed:10619841). Modulates transcription start site selection (PubMed:10318856). Exhibits also autoacetyltransferase activity that contributes to the activated transcription (PubMed:12931194).

Cellular Location

Nucleus. Chromosome. Note=Non-acetylated form colocalizes with DNA in the G0/1, S and G2 phases of the cell cycle, but not during mitosis (PubMed:24441171). Acetylated form colocalizes at transcriptionally silent mitotic chromatids during mitosis at metaphase, anaphase, and telophase phases of the cell cycle (PubMed:24441171).

Tissue Location

Expressed in the inner cell mass forming the embryoblast (PubMed:24441171). Not detected in cells from the outer thin layer trophoblast (at protein level) (PubMed:24441171)


GTF2B antibody - N-terminal region - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)

- [Flow Cytometry](#)
- [Cell Culture](#)

GTF2B antibody - N-terminal region - Images



WB Suggested Anti-GTF2B Antibody Titration: 0.2-1 µg/ml

ELISA Titer: 1:312500

Positive Control: Human brain

GTF2B antibody - N-terminal region - Background

General factor that plays a major role in the activation of eukaryotic genes transcribed by RNA polymerase II.

GTF2B antibody - N-terminal region - References

Ha I., et al. Nature 352:689-695(1991).
Malik S., et al. Proc. Natl. Acad. Sci. U.S.A. 88:9553-9557(1991).
Ota T., et al. Nat. Genet. 36:40-45(2004).
Goshima N., et al. Nat. Methods 5:1011-1017(2008).
Gregory S.G., et al. Nature 441:315-321(2006).