

**HCAR3 / GPR109B / HM74 Antibody (N-Terminus)**  
**Rabbit Polyclonal Antibody**  
**Catalog # ALS10059****Specification**

---

**HCAR3 / GPR109B / HM74 Antibody (N-Terminus) - Product Information**

|                   |                        |
|-------------------|------------------------|
| Application       | IHC                    |
| Primary Accession | <a href="#">P49019</a> |
| Reactivity        | Human                  |
| Host              | Rabbit                 |
| Clonality         | Polyclonal             |
| Calculated MW     | 44kDa KDa              |

**HCAR3 / GPR109B / HM74 Antibody (N-Terminus) - Additional Information****Gene ID** 8843**Other Names**

Hydroxycarboxylic acid receptor 3, G-protein coupled receptor 109B, G-protein coupled receptor HM74, G-protein coupled receptor HM74B, Niacin receptor 2, Nicotinic acid receptor 2, HCAR3, GPR109B, HCA3, HM74B, NIACR2

**Target/Specificity**

Human GPR109B / HM74. BLAST analysis of the peptide immunogen showed no homology with other human proteins, except HCAR2 (100%).

**Reconstitution & Storage**

Long term: -70°C; Short term: +4°C

**Precautions**

HCAR3 / GPR109B / HM74 Antibody (N-Terminus) is for research use only and not for use in diagnostic or therapeutic procedures.

**HCAR3 / GPR109B / HM74 Antibody (N-Terminus) - Protein Information****Name** HCAR3**Synonyms** GPR109B, HCA3, HM74B, NIACR2**Function**

Receptor for 3-OH-octanoid acid mediates a negative feedback regulation of adipocyte lipolysis to counteract prolipolytic influences under conditions of physiological or pathological increases in beta- oxidation rates. Acts as a low affinity receptor for nicotinic acid. This pharmacological effect requires nicotinic acid doses that are much higher than those provided by a normal diet.

**Cellular Location**

Cell membrane; Multi-pass membrane protein.

**Tissue Location**

Expression largely restricted to adipose tissue and spleen.


**Volume**

50  $\mu$ l

**HCAR3 / GPR109B / HM74 Antibody (N-Terminus) - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

**HCAR3 / GPR109B / HM74 Antibody (N-Terminus) - Images**

Anti-GPR109B / HM74 antibody IHC of human spleen.

**HCAR3 / GPR109B / HM74 Antibody (N-Terminus) - Background**

Receptor for 3-OH-octanoid acid mediates a negative feedback regulation of adipocyte lipolysis to counteract prolipolytic influences under conditions of physiological or pathological increases in beta-oxidation rates. Acts as a low affinity receptor for nicotinic acid. This pharmacological effect requires nicotinic acid doses that are much higher than those provided by a normal diet.

**HCAR3 / GPR109B / HM74 Antibody (N-Terminus) - References**

- Nomura H., et al. Int. Immunol. 5:1239-1249(1993).  
Suwa M., et al. Submitted (JUL-2001) to the EMBL/GenBank/DDBJ databases.  
Ota T., et al. Nat. Genet. 36:40-45(2004).  
Scherer S.E., et al. Nature 440:346-351(2006).  
Wise A., et al. J. Biol. Chem. 278:9869-9874(2003).