

CXCR2 Antibody (N-Terminus)
Rabbit Polyclonal Antibody
Catalog # ALS10169**Specification**

CXCR2 Antibody (N-Terminus) - Product Information

Application	IHC
Primary Accession	P25025
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal
Calculated MW	41kDa KDa

CXCR2 Antibody (N-Terminus) - Additional Information**Gene ID** 3579**Other Names**

C-X-C chemokine receptor type 2, CXC-R2, CXCR-2, CDw128b, GRO/MGSA receptor, High affinity interleukin-8 receptor B, IL-8R B, IL-8 receptor type 2, CD182, CXCR2, IL8RB

Target/Specificity

Human CXCR2. BLAST analysis of the peptide immunogen showed no homology with other human proteins, except EYA3 (50%).

Reconstitution & Storage

Long term: -70°C; Short term: +4°C

Precautions

CXCR2 Antibody (N-Terminus) is for research use only and not for use in diagnostic or therapeutic procedures.

CXCR2 Antibody (N-Terminus) - Protein Information**Name** CXCR2**Synonyms** IL8RB**Function**

Receptor for interleukin-8 which is a powerful neutrophil chemotactic factor (PubMed:1891716). Binding of IL-8 to the receptor causes activation of neutrophils. This response is mediated via a G- protein that activates a phosphatidylinositol-calcium second messenger system (PubMed:8662698). Binds to IL-8 with high affinity. Also binds with high affinity to CXCL3, GRO/MGSA and NAP-2.

Cellular Location

Cell membrane; Multi-pass membrane protein.


Volume
50 µl

CXCR2 Antibody (N-Terminus) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

CXCR2 Antibody (N-Terminus) - Images



Anti-CXCR2 antibody ALS10169 IHC of human kidney.

CXCR2 Antibody (N-Terminus) - Background

Receptor for interleukin-8 which is a powerful neutrophil chemotactic factor. Binding of IL-8 to the receptor causes activation of neutrophils. This response is mediated via a G-protein that activates a phosphatidylinositol-calcium second messenger system. Binds to IL-8 with high affinity. Also binds with high affinity to CXCL3, GRO/MGSA and NAP-2.

CXCR2 Antibody (N-Terminus) - References

Murphy P.M., et al. Science 253:1280-1283(1991).
Cerretti D.P., et al. Mol. Immunol. 30:359-367(1993).
Sprenger H., et al. J. Biol. Chem. 269:11065-11072(1994).
Ahuja S.K., et al. J. Biol. Chem. 269:26381-26389(1994).
Kato H., et al. Genes Immun. 1:330-337(2000).